

# Jahresbericht der Deutschen Mathematiker-Vereinigung

3 - 2003

Herausgegeben von A. Krieg unter Mitwirkung von U. Gather, E. Heintze, B. Kawohl, H. Lange, H. Triebel



Der "Jahresbericht" ist das offizielle Veröffentlichungsorgan der Deutschen Mathematiker-Vereinigung, für dessen inhaltliche Gestaltung im Auftrag des Präsidiums der jeweilige Herausgeber zuständig ist. Im "Jahresbericht" sollen vornehmlich Überblicksartikel über Teilgebiete der reinen und angewandten Mathematik, Nachrufe sowie historische Artikel und Buchbesprechungen veröffentlicht werden.

#### Manuskripte:

Alle für die Schriftleitung des Jahresberichts bestimmten Briefe und Manuskripte sind an Prof. Dr. A. Krieg zu richten. Für Buchbesprechungen ist Prof. Dr. H. Lange zuständig. Bücher, von denen eine Besprechung erfolgen soll, werden bei den Verlagen angefordert. Autoren von Buchbesprechungen und Artikeln werden gebeten, die vorhandenen LATEX-style-files für den Jahresbericht zu verwenden. Somit kann der Aufwand für die Satzarbeiten erheblich reduziert werden. Sollten Illustrationen in die Arbeiten integriert werden, können diese auch in das Satzsystem übernommen werden. Dazu ist es erforderlich, dass die Daten der Abbildungen nochmals in separaten Dateien einzeln abgespeichert werden. Die LATEX-style-files sind neben weiteren Informationen im Internet verfügbar unter

http://www.mathA.rwth-aachen.de/dmv/index.html

Auf Anfrage können die style-files auch auf Diskette zugeschickt werden.

Grundsätzlich sollen nur solche Manuskripte eingereicht werden, die nicht gleichzeitig an anderer Stelle zur Veröffentlichung eingereicht oder bereits veröffentlicht worden sind. Mit der Annahme zur Veröffentlichung erwirbt der Verlag das Recht zur Vervielfältigung und Verbreitung sowie das Recht der Übersetzung in andere Sprachen.

#### Verlag:

B. G. Teubner Verlag/GWV Fachverlage GmbH Abraham-Lincoln-Straße 46 65189 Wiesbaden http://www.teubner.de http://www.gwv-fachverlage.de

#### Abo-/Leserservice:

Tatjana Hellwig Telefon: (06 11) 78 78-1 51 Fax: (06 11) 78 78-4 23 E-Mail: tatjana.hellwig@bertelsmann.de Geschäftsführer: Dr. Hans-Dieter Haenel Verlagsleitung: Dr. Heinz Weinheimer Gesamtleitung Anzeigen: Thomas Werner Gesamtleitung Produktion: Reinhard van den Hövel Gesamtleitung Vertrieb: Gabriel Göttlinger

Jahresberich

der Deutschen Mathematiker-Vereinigung

#### Marketing/Sonderdrucke:

Stefanie Hoffmann Telefon: (06 11) 78 78-3 79 Fax: (06 11) 78 78-4 39 E-Mail: stefanie.hoffmann@bertelsmann.de

#### Abonnentenverwaltung:

(Änderungen von Adressen und Bankverbindung, Rückfragen zu Rechnungen oder Mahnung) VVA-Zeitschriftenservice, Abt. D6F6 / Jahresbericht der Deutschen Mathematiker-Vereinigung, Postfach 7777, 33310 Gütersloh Ursula Müller Telefon: (0 52 41) 80-19 65 Fax: (0 52 41) 80-96 20 E-Mail: ursula.mueller@bertelsmann.de

#### **Bezugsbedingungen:**

Die Zeitschrift erscheint 4mal jährlich zum Jahresabonnementpreis von € 98,- (176,- sFr) inkl. Versandkosten. Der Bezug von Einzelheften ist nicht möglich. Schriftliche Kündigung des Abonnements spätestens sechs Wochen vor Ablauf des Bezugsjahres.

Für persönliche Mitglieder der DMV, die den Jahresbericht zu beziehen wünschen, ist der zwischen DMV und Verlag vereinbarte Bezugspreis maßgebend, der im Rahmen des Mitgliedsbeitrags erhoben wird.

#### Copyright ©

B. G. Teubner Verlag/GWV Fachverlage GmbH, Wiesbaden 2003. Printed in Germany. Der Verlag B. G. Teubner ist ein Unternehmen der Fachverlagsgruppe BertelsmannSpringer. Alle Rechte vorbehalten. Kein Teil dieser Zeitschrift darf ohne schriftliche Genehmigung des Verlages vervielfältigt oder verbreitet werden. Unter dieses Verbot fällt insbesondere die gewerbliche Vervielfältigung per Kopie, die Aufnahme in elektronischen Datenbanken und die Vervielfältigung auf CD-ROM und allen anderen elektronischen Datenträgern.

Satz: Fotosatz Behrens, D-68723 Oftersheim Druck: Thomas Müntzer, Bad Langensalza

ISSN 0012-0456

## Inhalt

## Jahresbericht der Deutschen Mathematiker-Vereinigung, 105. Bd. 2003, Nr. 3

Übersichtsartikel	Historischer Artikel	Buchbesprechungen
-------------------	----------------------	-------------------

Jahresbericht der Deutschen Mathematiker-Vereinigung, 105. Bd. 2003, Nr. 3

#### In den nächsten Heften erscheinende Arbeiten

G. Hiss: Die Sporadischen Gruppen

**D. Horstmann:** From 1970 to present: The Keller-Segel model in chemotaxis and consequences II

K. Mohnke: Holomorphe Kurven, Hamiltonsche Dynamik und Symplektische Topologie

I. Pigeot, K. Bammann, A. Reineke, N. Wawro, A. Zierer: Statistical methods in genetics: From microarrays to genetic epidemiology – an overview

#### Anschriften der Herausgeber

Prof. Dr. Aloys Krieg,	Lehrstuhl A für l	Mathematik,	RWTH	Aachen,	52056	Aachen
E-Mail: krieg@mat	hA.rwth-aachen	.de				

Prof. Dr. Ursula Gather, Lehrstuhl für Mathematische Statistik und industrielle Anwendungen, Universität Dortmund, 44221 Dortmund E-Mail: gather@statistik.uni-dortmund.de

Prof. Dr. Ernst Heintze, Institut für Mathematik, Universität Augsburg, 86135 Augsburg E-Mail: heintze@math.uni-augsburg.de

Prof. Dr. Bernhard Kawohl, Mathematisches Institut, Universität zu Köln, 50923 Köln E-Mail: kawohl@mi.uni-koeln.de

Prof. Dr. Herbert Lange, Mathematisches Institut, Friedrich-Alexander-Universität Erlangen-Nürnberg, Bismarckstraße 1<sup>1</sup>/<sub>2</sub>, 91054 Erlangen E-Mail: lange@mi.uni-erlangen.de

Prof. Dr. Hans Triebel, Mathematisches Institut, Friedrich-Schiller-Universität, Ernst-Abbe-Platz 1–4, 07740 Jena E-Mail: triebel@minet.uni-jena.de

#### Bezugshinweis

Früher erschienene Bände (ab Band 68) des "Jahresberichts der Deutschen Mathematiker-Vereinigung" können durch den Buchhandel oder den Verlag bezogen werden.

Nachdruck der Bände 41 bis 67 liefert: Swets & Zeitlinger, Heereweg 347 b, POB 810, NL-2160 SZ Lisse/Holland

Jahresbericht der Deutschen Mathematiker-Vereinigung, 105. Bd. 2003, Nr. 3

## Vorwort

Zu den vom DMV-Präsidium beschlossenen Neuerungen beim Jahresbericht gehört auch die Möglichkeit, Nachwuchswissenschaftlern eine Chance zu bieten, sich einer breiten Öffentlichkeit zu präsentieren. In diesem Heft finden Sie den Übersichtsartikel von D. Horstmann über das Keller-Segel Modell in der Chemotaxis. Darüber hinaus kommen wir damit dem Wunsch vieler Leser nach, anwendungsbezogene Beiträge der Mathematik zu publizieren, die in diesem Fall tiefliegende Methoden der Analysis verwenden.

Aufgrund der Umfangsbeschränkungen können in diesem Heft keine Buchbesprechungen erscheinen, denen aber in Heft 4 ein größerer Anteil eingeräumt werden wird.

A. Krieg





## From 1970 until present: The Keller-Segel model in chemotaxis and its consequences I

**Dirk Horstmann** 

## Abstract

- Keywords and Phrases: Chemotaxis equations, steady state analysis, global existence, finite time blow-up, invariant sets
- Mathematical Subject Classification: 35B30, 35J20, 35J25, 35J65, 35K50, 35K57, 92C17

This article summarizes various aspects and results for some general formulations of the classical chemotaxis models also known as Keller-Segel models. It is intended as a survey of results for the most common formulation of this classical model for positive chemotactical movement and offers possible generalizations of these results to more universal models. Furthermore it collects open questions and outlines mathematical progress in the study of the Keller-Segel model since the first presentation of the equations in 1970.

Eingegangen: 24.03.2003

Dirk Horstmann, Mathematisches Institut der Universität zu Köln, Weyertal 86–90, D-50931 Köln, Germany. Email: dhorst@mi.uni-koeln.de DMV JAHRESBERICHT DER DMV © B. G. Teubner 2003

## 1 Introduction

Mathematical analysis of biological phenomena has become more and more important in understanding these complex processes. Thus, the number of mathematicians studying biological and medical phenomena and problems is continuously increasing in recent years. One such biological topic is the movement of population densities or the movement of single particles. Changes in the environment of mobile species can influence its movement. For example, humans sense their environment and given a particular situation or the state of the environment, they make their decisions as to where to move. For example we might be attracted by a tantalizing smell and move towards it, since we expect a delicious food, or we move away from a place if there is a repellent odor. Animals and humans also use this effects (for example) to attract mating partners with special colorful feathers or with enticing perfumes etc. In [89, 90] one can find the silk moth *Bombyx mori* as an example of a species that uses a special odor to attract a mating partner. During mating season the female moth secretes a scent caused by a pheromone bombykol which attracts the male to move in direction of the increasing concentration of this scent. This helps the male moths to find the female. Before presenting another example where changes in the environment affect the movement of a mobile species let me cite the following anecdote from the German news magazine "DER SPIEGEL" 36/1998 [46] that is said to have happened in the late 1950s at Princeton University:

"The genius was stunned. In the late 1950s Albert Einstein watched disbelievingly a film of the young scientist John Tyler Bonner at Princeton University. The star of the movie was an unimpressive tiny creature: an amoeba called Dictyostelium.(...) As soon as "Dicty",(...), starts to become hungry it undergoes a miraculous metamorphosis.(...) The Dictys become one.(...) Einstein's question is still unsolved: Why does Dicty undergo a deadly intermezzo as a complex multicellular organism to live then alone and autistically?" (Quoted from [46] translated by the author.)

The cellular slime mold Dictyostelium discoideum was discovered by K. B. Raper in 1935 and in the subsequent years aroused the interest of many scientists. Nowadays Dictyostelium discoideum is a model organism for biomedical research of the National Institutes of Health (NIH). One reason for the growing interest in this cellular slime mold was caused by the fact that "development in Dictyostelium discoideum results only in two terminal cell types, but processes of morphogenesis and pattern formation occur as in many higher organisms" (see [107, page 354]). This raised the hope of biologists that studying this cellular slime mold might aid in the understanding of the secret of cell differentiation. But what initiates the change from a single cell organism to a complex multicellular organism? And how does this process take place?

During its life cycle a myxamoebae population of the Dictyostelium grows by cell division as long as there is sufficient nourishment. When the food resources are exhausted the myxamoebae spread over the entire domain available to them. After a while one cell starts to exude cyclic Adenosine Monophosphate (cAMP) which attracts the other myxamoebae. The myxamoebae begin to move towards the so-called founder cell and are also stimulated to emit cAMP. The myxamoebae aggregate and start to differentiate. At the end of the aggregation the myxamoebae form a pseudoplasmoid, in which every myxamoebae maintains its individual integrity. This pseudoplasmoid moves towards light sources. After a time a fruiting body is formed and spores are spread. Thus the life cycle begins again. For more details on the life cycle of the Dictyostelium I refer to [15], for example.

A reaction to an external stimulus is generally called taxis and is then specified by describing the reason for the reaction. Therefore, there are many different tactical responses such as chemotaxis, galvanotaxis and phototaxis. In this article I will focus on chemotactical movement of mobile species which can lead to various different pattern formations.

Chemotaxis is the influence of chemical substances in the environment on the movement of mobile species. This can lead to strictly oriented movement or to partially oriented and partially tumbling movement. The movement towards a higher concentration of the chemical substance is termed positive chemotaxis and the movement towards regions of lower chemical concentration is called negative chemotactical movement. Thus, the Bombyx mori and the Dictyostelium discoideum are two species that move in a chemotactically positive manner towards the higher concentration of the bombykol resp. the cAMP. The substances that lead to positive chemotaxis are chemoattractants and those leading to negative chemotaxis are so-called repellents.

Chemotaxis is an important means for cellular communication. Communication by chemical signals determines how cells arrange and organize themselves, like for instance in development or in living tissues. A large number of examples for mobile species behaving in a chemotactical manner are known. In addition to the above mentioned two examples, I would also like to draw attention to a third species, the bacterial strain *Rhizobia meliloti*. As described in [38], this bacterial strain responds chemotactically to root exudates isolated from the soil of leguminous plants. The bacterial strain in the surrounding soil of the plants are guided to nodules in the roots of nitrogen-fixing plants by a chemical gradient. Therefore, they play an important role in agricultural ecology.

One aspect during positive oriented chemotactical movement is the formation of cells (amoebae, bacteria, etc) amounts during the responds of the species population to the change of the chemical concentrations in the environment. Such aggregation patterns often require a certain threshold number of individuals. Therefore, depending on the case in question, that is the species being observed, such threshold phenomena should be reflected in the model. For example aggregation in Dictyostelium is only possible if the total number of myxamoebae in the population is larger than a threshold number of myxamoebae. In [26] the threshold value of  $5 \cdot 10^4$  myxamoebae per  $cm^2$  is given for Dictyostelium discoideum. This chemotactical effect has been observed in experiments to demonstrate chemotaxis of bacteria (see for example [120]). Positive and negative chemotaxis can be studied in petri dish cultures. If the bacteria are placed in the center of the dish of agar that contains an attractant, the bacteria will exhaust the local supply and then move outward following the attractant gradient they have created. This results in an expanding ring of bacteria. To demonstrate negative chemotaxis one can place a disk of repellent in a petri dish of semisolid agar and bacteria. The bacteria will then move away from the repellent. This movement away from the repellent will lead to the creation of a clear zone around the disk.

Übersichtsartikel	
-------------------	--

Alternatively, one can demonstrate chemotaxis by observing bacteria in the chemical gradient produced when a thin capillary tube is filled with an attractant and lowered into a bacterial suspension. While the attractant diffuses, the bacteria collect and move up the tube. The observed positive chemotactical effect in this experiment is the formation of bacteria (myxamoebae, cells, etc.) bands. Such and similar experiments have been carried out for example by Adler [1]. Adler's observations correspond to the formation of traveling waves and pulses that spread through the population density. Thus an interesting question is whether or not the mathematical models describing chemotactical movement have traveling wave solutions.

These phenomena have motivated a large number of scientists to study chemotaxis and to use the mathematical language to describe the observed phenomena. The intention of the present survey is to collect the results for a classical model describing chemotactical movement, to expose the lines of research.

The outline of the first part of this survey is as follows:

In the second section two different approaches for modeling chemotaxis will be inculded. This section will also introduce the "classical" chemotaxis model by Keller and Segel, the center of our considerations for the remainder of the paper. The third section is devoted to steady-state analysis for this classical model by Keller and Segel done so far. It will be shown that all the effects demonstrated in the analysis depend on the functional forms of the three main processes during chemotactical movement. They are:

- a) The sensing of the chemoattactant which has an effect on the oriented movement of the species.
- b) The production of the chemoattractant by a mobile species or an external source.
- c) The degradation of the chemoattractant by a mobile species or an external effect.

Within the context of steady-state analysis the focus will be on a linear chemotactic sensitivity function but will also collect the results for different versions of the Keller-Segel model. When appropriate I will summarize results for other sensitivity functions in a table at the end of a section. Section 4 will deal with the possibility of an explosion of the solution in finite time in the case of a linear sensitivity function. Here I point out the different lines of research in chronological order. This can be accomplished without losing clarity in the results. Section 5 addresses questions asked in [69] on the possibility of explosion of the solution in finite time in the case of a linear sensitivity function. This section is then followed by generalizations of these results for other more general versions of the classical model in Section 6. In the seventh section of the present article I will present some comparison results for some general versions of the Keller-Segel model proved by Wolfgang Alt in his Habilitation [3]. This section, however, will be somewhat technical. At some places in the text questions will be formulated that arise from the results stated in the article. These questions are partially answered in subsequent sections. but some are still open problems and might be worth further study. I will close this first part of the summary of results for the Keller-Segel model with some brief comments on other approaches and models for chemotaxis.

In the second part of this survey [67a] I will turn to special solutions for the Keller-Segel model in chemotaxis. It will summarize the results on self-similar and traveling wave solutions for this classical chemotaxis model. The outline of [67a] is as follows:

After a short introduction I will turn to the results on self-similar solutions. Then I will go a bit more into detail on traveling wave solutions for Keller-Segel type systems. Known results will be presented and discussed. Furthermore I will discuss which functional forms for sensing, producing and degrading the chemoattractant might cause resp. favor the existence of traveling wave solutions. I will also sketch some results on traveling front solutions for a chemotaxis system with population growth. Finally I will close this summary of results for the Keller-Segel model with some brief comments.

## 2 Different perspectives to model chemotactical movement and the formulation of the classical chemotaxis equations

Modeling chemotactical movement of mobile species can be done from two different perspectives: either from the microscopic or from the macroscopic perspective. Both approaches have been used over years and the derivation of the macroscopic equations from the microscopic, or to be precise the validation of the passage to the limiting equations is still a topic that is studied by a large number of scientists and depending on the model is still an open problem.

## 2.1 The macroscopic perspective

The first approach that should be presented here is the macroscopic perspective where one considers the whole population respectively the density of the population at one place and one time directly. This approach leads to a continuous reaction-diffusion model where the diffusion of the population density is modeled with Fourier's and Fick's laws and in which the reactions are viewed as functions of the population density and possibly some external signal or control substance.

In the year 1970 Evelyn Fox Keller and Lee A. Segel used this perspective to present a system of four strongly coupled parabolic partial differential equations, which describes the aggregation of cellular slime molds like the Dictyostelium discoideum. Let u(t, x) denote the myxamoebae density of the cellular slime molds and v(t, x) denote a chemoattractant concentration at time t in point x. To model the aggregation of a cellular slime mold population they assume in [71] the following basic processes that take place during the aggregation phase:

- (a) The chemoattractant is produced per amoeba at a rate f(v).
- (b) There exist an extracellular enzyme that degrades the chemoattractant. The concentration of the is enzyme at time t in point x is denoted by p(t, x). This enzyme is produced by the myxamoebae at a rate g(c, p) per amoeba.

	10	Contraction of the second	
IIh	orcic	htoo	rtillal
UD	CISIC	JIIISa	IUNCI

(c) The chemoattractant and the enzyme react to form a complex  $\mathcal{E}$  of concentration  $\eta$  which dissociates into a free enzyme plus the degraded product.

$$v + p \stackrel{r_1}{\xrightarrow{}} \mathcal{E} \stackrel{r_2}{\xrightarrow{}} p + \text{ degraded product}$$

(d) The chemoattractant, the enzyme and the complex diffuse according to Fick's law.

The balance of the myxamoebae density u(t, x) in any control volume D (which holds for example in the special case of Dictyostelium discoideum aggregation) implies the equation

(1) 
$$\frac{d}{dt}\int_{D}u(t,x) dx = -\int_{\partial D}(J^{(u)}(t,x)\cdot n(x))dS.$$

Here  $J^{(u)}(t, x)$  denotes the flow of the myxamoebae density. This flow contains according to Fick's law a part that is proportional to the density gradient and according to Fourier's law for the heat flow a part that is proportional to the chemoattractant gradient. Thus we see that:  $J^{(u)}(t, x) = k_2 \nabla v - k_1 \nabla u$ . As a chemical substance the chemoattractant diffuses and we get

(2) 
$$\frac{d}{dt}\int_{D}v(t,x)\ dx = Q^{(v)}(t,D) - \int_{\partial D}(J^{(v)}(t,x)\cdot n(x))dS,$$

where  $Q^{(\nu)}(t,D)$  denotes the produced chemoattractant  $\nu(t,x)$  per domain and time volume. The flow  $J^{(\nu)}(t,x)$  is given by:  $J^{(\nu)}(t,x) = -k_c \nabla \nu$ . Assuming the analogous equations for the enzyme and the complex, and taking the basic processes into account we derive at the following system:

$$(3) \begin{cases} u_t = \nabla(k_1(u,v)\nabla u - k_2(u,v)\nabla v), & x \in \Omega, \ t > 0\\ v_t = k_c\Delta v - r_1vp + r_{-1}\eta + uf(v), & x \in \Omega, \ t > 0\\ p_t = k_p\Delta p - r_1vp + (r_{-1} + r_2)\eta + ug(v,p), & x \in \Omega, \ t > 0\\ \eta_t = k_\eta\Delta \eta + r_1vp - (r_{-1} + r_2)\eta, & x \in \Omega, \ t > 0\\ \partial u/\partial n = \partial v/\partial n = \partial p/\partial n = \partial \eta/\partial n = 0, & x \in \partial \Omega, \ t > 0\\ u(0,x) = u_0(x), \ v(0,x) = v_0(x), & x \in \Omega, \\ p(0,x) = p_0(x), \ \eta(0,x) = \eta_0(x), & x \in \Omega, \end{cases}$$

where  $r_{-1}$ ,  $r_1$  and  $r_2$  are constants representing the reaction rates mentioned in assumption (c). Here  $\Omega$  denotes a bounded domain in  $\mathbb{R}^N$  with boundary  $\partial \Omega$ .

Let us simplify the chemical processes in the life cycle via assuming that the complex is in a steady state with regard to the chemical reaction and that the total concentration of the free and the bounded enzyme is a constant. Thus one gets a simplified formulation of this original Keller-Segel model that has already been proposed by E. F. Keller and L. A. Segel themselves to reduce their original system of four strongly coupled parabolic equations to a model that is as simple as possible. Thus their motto that "*it is useful for the sake of clarity to employ the simplest reasonable model*" (see [71, page 403]) leads them to the following system of "only two" strongly coupled nonlinear parabolic equations:

(4) 
$$\begin{cases} u_t = \nabla(k_1(u,v)\nabla u - k_2(u,v)\nabla v), & x \in \Omega, \ t > 0\\ v_t = k_c \Delta v - k_3(v)v + uf(v), & x \in \Omega, \ t > 0\\ \partial u/\partial n = \partial v/\partial n = 0, & x \in \partial \Omega, \ t > 0\\ u(0,x) = u_0(x), \ v(0,x) = v_0(x), & x \in \Omega. \end{cases}$$

However, it might be necessary to remember the original system if one tries to describe certain pattern formations during the aggregation of some particular species. It is possible that the reduction to two equations that was done in [71] was too restrictive to cover all observable generated patterns during the aggregation of mobile species.

## 2.2 The microscopic approach

From the microscopic perspective one interprets the movement of species populations as a consequence of microscopic irregular movement of single members of the considered population that results in a macroscopic regular behaviour of the whole population. This then leads in a parabolic limit to reaction-diffusion processes, however, in this case the passage to the continuum limit of the microscopic problem and thus studying the resulting, continuous partial differential equations has to be valid and justified. Usually it is assumed that in a particles population each single particle moves around in a random walk. Leaving the justification of the limiting process open, this approach gives us at least a formal way to derive reaction-diffusion processes from the microscopic point of view.

For example in [116] H. G. Othmer and A. Stevens used the microscopic perspective and started with a continuous-time, discrete-space random walk for a *single particle* in one space dimension. Restricting themselves to one step jumps and assuming that the conditional probability  $p_i(t)$  that a walker is in  $i \in \mathbb{Z}$  at time t – conditioned on the fact that it begins in i = 0 at t = 0 – evolves according to the continuous time master equation

(5) 
$$\frac{\partial p_i}{\partial t} = \mathcal{T}^+_{i-1}(W) \ p_{i-1} + \mathcal{T}^-_{i+1}(W) \ p_{i+1} - (\mathcal{T}^+_i(W) + \mathcal{T}^-_i(W)) \ p_i.$$

Here  $\mathcal{T}_i^{\pm}(\cdot)$  are the transition probabilities per unit time for a one-step jump to  $i \pm 1$ , and  $(\mathcal{T}_i^+(W) + \mathcal{T}_i^-(W))^{-1}$  is the mean waiting time at the *i*<sup>th</sup> site. It is assumed that these are nonnegative and suitably smooth functions of their arguments. The vector Wis given by  $W = (\cdots, w_{-i-1/2}, w_{-i}, w_{-i+1/2}, \cdots, w_o, w_{1/2}, \cdots)$ . For generality and in context with a self-attracting reinforced random walk analyzed by Davis [28] the density of the control species w is defined on the embedded lattice of half the step size. As (5) is written, the jump probabilities may depend on the entire state and on the entire distribution of the control species. Since there is no explicit dependence on the previous state the process may appear to be Markovian, but if the evolution of  $w_i$  depends on  $p_i$  then there is an implicit history dependence, and thus the jump process by itself is not Markovian. However, the composite process for the evolution (p, w) is a Markov process. There are three distinct types of models that are considered in [116], which differ in the dependence of the transition rates on  $w_i$ : (i) strictly local models in which for example

Ubersichtsartikel	Historischer Artikel	Buchbesprechungen

 $\mathcal{T}_i^{\pm}$  are equal, (ii) barrier models in which for example  $\mathcal{T}_i^{\pm}(W) = \mathcal{T}_i(w_{i\pm 1/2})$ , and (iii) gradient models for example with  $\mathcal{T}_{i-1}^+(W) = \alpha + \beta(\tau(w_i) - \tau(w_{i-1}))$  and  $\mathcal{T}_{i+1}^-(W) = \alpha + \beta(\tau(w_i) - \tau(w_{i+1}))$  for  $\alpha \ge 0$  and a function  $\tau$  of the control substance.

Considering a grid of mesh size h and setting x = ih the *formal* expansion of the righthand side of equation (5) as a function of x to second order in h leads

1. in case (i) to

$$\frac{\partial p}{\partial t} = h^2 \frac{\partial^2}{\partial x^2} (\mathcal{T}(w)p) + \mathcal{O}(h^4)$$

and so with an assumed scaling  $\lim_{h\to 0,\lambda\to 0} \lambda h^2 = D$ , where  $\lambda$  has dimension  $t^{-1}$ , to the limiting problem

$$\frac{\partial p}{\partial t} = D \frac{\partial^2}{\partial x^2} (\mathcal{T}(w)p),$$

2. in case (ii) with the same scaling to

$$\frac{\partial p}{\partial t} = D \frac{\partial}{\partial x} \left( \mathcal{T}(w) \frac{\partial p}{\partial x} \right),$$

3. and in case (iii) once again with the same scaling as before to

$$\frac{\partial p}{\partial t} = D \frac{\partial}{\partial x} \left[ p \left( \frac{\alpha}{p} \frac{\partial p}{\partial x} - 2\beta \tau'(w) \frac{\partial w}{\partial x} \right) \right].$$

Assuming various possible evolution equations of the control substance *w* this leads formally from a random walk of a single particle to a limiting diffusion equation *for the probability of one particle to be located in x at time t*. Of course one can also study these equations as an ad hoc approach for particle densities, but their derivations are then only formal approaches and by far not rigorous. The key problem to derive the limiting equations from the multi particle random walk is the interaction of the particles via the control species. A rigorous derivation of limiting equations in these cases is not done yet. Simulations of these are presented in [116] and [141].

The first rigorous derivation of chemotaxis equations from a microscopic model, namely an interacting stochastic many-particle system, has been done in [139] and [143]. In [143] Stevens proved that for large enough particle numbers the dynamics of the below given interacting particle systems are well described by the solution of chemotaxis systems which for this case describe population densities. Explicit error estimates are also given. For the derivation it was assumed that every particle interacts mainly with those of the other particles which are located in a certain neighbourhood of itself. The neighbourhood is macroscopically small and microscopically large. As a consequence the interaction range between the particles is shrinking as the number of particles goes to  $\infty$ , while the number of particles in the shrinking neighbourhoods is also growing to  $\infty$ .

So let the subscript u mark the terms related to bacteria and the subscript v mark the terms related to the chemical substance slime particles. Let  $S(M,t) = S_u(M,t) + S_v(M,t)$  denote the set of all particles in a *M*-particle system, the particles are numbered consecutively by taking a new number for each new-born particle.  $P_M^k(t) \in \mathbb{R}^d$ ,

 $k \in S(M, t)$  describes the position of the *k*th particle at time  $t \ge 0$ . Furthermore let  $\delta_x$  denote the Dirac measure at  $x \in \mathbb{R}^d$ ,  $d \in \mathbb{N}$ . Stevens considered the following measure valued empirical processes:

$$t \mapsto S_{M_{\mathcal{U}}}(t) = \frac{1}{M} \sum_{k \in S_{\mathcal{U}}(M,t)} \delta_{P_{M}^{k}(t)},$$
$$t \mapsto S_{M_{\mathcal{V}}}(t) = \frac{1}{M} \sum_{k \in S_{\mathcal{V}}(M,t)} \delta_{P_{M}^{k}(t)} \text{ and }$$
$$S_{M}(t) = S_{M_{\mathcal{U}}}(t) + S_{M_{\mathcal{V}}}(t).$$

The dynamics of the particles depend on the following smoothed versions of  $S_{M_{\mu}}$ ,  $S_{M_{\nu}}$ :

$$\hat{s}_{M_{u}} = \left(S_{M_{u}} * W_{M} * \hat{W}_{M}\right)(x), \ \hat{s}_{M_{v}} = \left(S_{M_{v}} * W_{M} * \hat{W}_{M}\right)(x)$$

where  $W_M$  and  $\hat{W}_M$  are moderately scaled functions of a fixed symmetric function  $W_1$ (e.g. a Gaussian):  $W_M = M^{\alpha} W_1(M^{\alpha/d}x)$  and  $\hat{W}_M = M^{\hat{\alpha}} W_1(M^{\hat{\alpha}/d}x)$ , where  $\alpha$  and  $\hat{\alpha}$  are constants that for technical reasons have to fulfill certain smallness conditions (see [143, page 4]). Setting up the corresponding Fokker-Planck equations for each particle and taking the particle interaction into account she ends up with the following equation

(6) 
$$dP_{M}^{k}(t) = \chi_{M}(t, P_{M}^{k}(t)) \nabla \hat{s}_{M_{v}}(t, P_{M}^{k}(t)) dt + \sqrt{2\mu} dW^{k}(t),$$

where  $W^k(\cdot)$  are independent  $\mathbb{R}^d$ -valued standard Brownian motions,  $\mu > 0$  is a constant and  $\chi_M(t,x)$  is given by the equation  $\chi_M(t,x) = \chi(\hat{s}_{M_u}(t,x), \hat{s}_{M_v}(t,x))$  with a smooth function  $\chi : \mathbb{R}^+ \times \mathbb{R}^+ \to \mathbb{R}^+$ .

Under some technical assumptions she ends up with the weak formulation of the classical chemotaxis system

$$\begin{cases} u_t = \nabla(\mu \nabla u - \chi(u, v)u \nabla v) \\ v_t = \eta \Delta v - \gamma(u, v)v + \beta(u, v)u \end{cases}$$

where  $S_{M_u} \to u$ ,  $S_{M_v} \to v$  as  $M \to \infty$  in probability,  $\eta > 0$  is a constant and  $\gamma(\cdot, \cdot)$  and  $\beta(\cdot, \cdot)$  are smooth, positive functions on  $\mathbb{R}^+ \times \mathbb{R}^+$ . The derivation of the limit dynamics is done by extensions of techniques used by Oelschläger [112]. For further results on these aspects we refer the interested reader to [116, 142, 139, 140] and [143].

Another paper that should be mentioned in the context of the derivation of the Keller-Segel equations as a model for population densities from the kinetic equations is [4]. Denoting the density of individuals moving at (t, x) in direction  $\theta$  and having started their run at time  $\tau$  by a smooth function  $\sigma(t, x, \theta, \tau)$  W. Alt started in [4] with the differential-integral system

(7) 
$$\frac{\partial}{\partial t}\sigma(t,x,\theta,\tau) + \frac{\partial}{\partial \tau}\sigma(t,x,\theta,\tau) + \theta\nabla_x(c(t,x)\sigma(t,x,\theta,\tau)) = -\beta(t,x,\theta,\tau)\sigma(t,x,\theta,\tau)$$

for  $\tau > 0$ ,  $\theta \in S^{n-1}$  (= the unit sphere in *n*-dimensional space), and speed c(t, x) of an individuum from the beginning of the run that stops at time *t* and point *x*, with a given probability  $\beta(t, x, \theta, \tau)$  per unit time. Here we have that

$$\sigma(t, x, \eta, 0) = \int_{0}^{\infty} \int_{S^{N-1}} \beta(t, x, \theta, \tau) \sigma(t, x, \theta, \tau) k(t, x, \theta, \eta) \ d\theta \ d\tau$$

for each  $\eta \in S^{N-1}$ , where  $k(t, x, \theta, \eta)$  denotes the given probability of a new chosen direction  $\eta$  after an individuum has stopped a run with direction  $\theta$  at (t, x). W. Alt shows that under some additional boundedness assumptions and hypotheses relating the size of some appearing parameters the density

$$\overline{u}(t,x) := \int_{S^{N-1}} \overline{\sigma}(t,x,\theta) \ d\theta = \int_{S^{N-1}} \left( \int_{0}^{\infty} \sigma(t,x,\theta,\tau) \ d\tau \right) \ d\theta$$

satisfies the first equation of the Keller-Segel model.

Last but not least I should mention the results from C. S. Patlak [118] in this section. In his paper from 1953 C. S. Patlak derives the partial differential equation of the random walk problem with persistence of direction and external bias. Here persistence of direction or internal bias means that the probability a particle travels in a given direction is not necessarily the same for all directions, but depends only on the particle's previous direction of motion. External bias means that the probability a particle travels in a given direction is dependent upon an external force on the particle. However, instead of speaking of the probability that a particle is at a point, Patlak speaks of a large number of particles moving around and therefore of the density of the particles about a point as a measure of the required probability. Thus in his picture of a random walk he speaks of a particle traveling in a straight line for a certain length of time  $\tau$  with an average speed c before turning, where the turning means a change in direction of the particle's motion. To make the idea of a random walk completely explicit – as opposed to diffusion – Patlak assumes that the particles have negligible interactions with each other and thus collision between the particles can be ignored. So let me list up the assumptions that Patlak uses throughout his paper:

- 1. The particles have negligible interaction with each other.
- 2. Each time the particle turns it start off afresh, with no "memory" of its previous c and  $\tau$ .
- 3. The amount of time spent in turning is negligible compared to the time the particle spends traveling between turns.
- 4. During a unit length of time the number of particles in each small reagion, as well as the distributions of c and  $\tau$ , remain approximately the same.

For the net displacement of a particle Patlak assumes that the probability of travel in any direction after turning and the distance of travel in a given direction are not necessarily the same for all directions. Now using the assumptions above he derives a modified Fokker-Planck equation. From this the partial differential equation for the random walk with persistence and external bias is obtained, which is more or less the first equation of the Keller-Segel model. Even though these results by Patlak are older than the paper by Keller and Segel system (4) is known as "the classical chemotaxis model" resp. as "the Keller-Segel model in chemotaxis".

Since it is not the goal of the present paper to go into the precise details of the derivations and approaches of [4, 116, 118, 139] and [142] we now leave this topic of the different possibilities to model chemotaxis and move to the main goal of the present paper, namely a review of the achieved results for system (4) which – as we have seen – can be derived from the macroscopic and microscopic perspective on chemotactical movement.

## 3 Linear stability analysis for the uniform distribution and nonconstant steady state solutions

Studying the steady state problem of the model (4) is already a challenging mathematical problem, showing a large variety of interesting aspects and uses a lot of astute mathematical techniques. Some tools used for the steady state analysis for the Keller-Segel model performed until now were techniques from the calculus of variations to show the existence of nonconstant stationary solutions and the existence of spike solutions. One tool used in this context is for example the mountain pass theorem by Ambrosetti and Rabinowitz [146, Theorem 6.1., page 109]. But let us proceed step by step to illustrate the way of progress on this topic.

In their paper from 1970 E. F. Keller and L. A. Segel studied in the case of two spatial dimensions the stability of a uniform state  $(u_0, v_0)$  for the species and the chemical attractant. Studying the effect of small (time dependent) perturbations of these uniform distributions they found by Taylor expansions in u and v of the right hand sides of the equations in (4) around the uniform state the following instability condition. The uniform distribution is unstable if

$$\frac{k_2(u_0,v_0)f(v_0)}{k_1(u_0,v_0)(k_3(v_0)+v_0k_3'(v_0))} + \frac{u_0f'(v_0)}{k_3(v_0)+v_0k_3'(v_0)} > 1,$$

or equivalent if

(8) 
$$\frac{k_2(u_0, v_0)v_0}{k_1(u_0, v_0)u_0} + \frac{u_0 f'(v_0)}{k_3(v_0) + v_0 k'_3(v_0)} > 1,$$

since a uniform state  $(u_0, v_0)$  satisfies the equality  $u_0 f(v_0) = v_0 k_3(v_0)$ . Here Keller and Segel call the uniform solution stable if the time dependent perturbations of the uniform distribution decrease with time. On the other hand they call the uniform distribution unstable, if these perturbations lead to solutions of (4) that increase in time.

Even though Keller's and Segel's stability analysis of the uniform state in [71] and the presented instability criterion is valid for a very general formulation of the system, the next "landmark" in the studies of the Keller-Segel model was the paper by V. Nanjundiah [106]. In that paper Nanjundiah performs a non-linear stability analysis for some versions of the Keller-Segel model in space dimension N = 2. In a linear stability analysis he first re-derives the instability criterion for the uniform distribution of myxamoebae and cAMP. Then his non-linear stability analysis for the system given by the equations

$$\begin{cases} u_t = \nabla(\nabla u - u\nabla\Phi(v)), & x \in \Omega, \ t > 0\\ v_t = k_c\Delta v - \gamma v + \alpha u, & x \in \Omega, \ t > 0\\ \partial u/\partial n = \partial v/\partial n = 0, & x \in \partial\Omega, \ t > 0\\ u(0, x) = u_0(x), \ v(0, x) = v_0(x), & x \in \Omega, \end{cases}$$

where  $k_c$ ,  $\gamma$ ,  $\alpha$  are positive constants strictly larger than zero and  $\Phi(v)$  denoting a che-

Übersichtsartikel	Historischer Artikel

motactical sensitivity function, leads him to one key statement that is mentioned in [106, page 102] for the case of a linear or a logarithmic chemotactical sensitivity function  $\Phi(v)$ . He states the following:

"The end-point (in time) of the aggregation is such that the cells are distributed in the form of  $\delta$ -function concentrations."

We want to sketch Nanjundiah's arguments leading to the above expectation for the case of a linear sensitivity function. So V. Nanjundiah considered in this case the following steady state system:

(9) 
$$\begin{cases} 0 = \nabla(\nabla u - u\nabla v), & x \in \Omega, \\ 0 = \Delta v - v + u, & x \in \Omega, \\ 0 = \partial u/\partial n = \partial v/\partial n, & x \in \partial \Omega. \end{cases}$$

For a general solution (u, v) of this system we see that the mean values over  $\Omega$  of u and v are equal to the same constant. The first equation of (9) motivates us to define a new function  $\psi(x)$  by  $u(x) = \psi(x)e^{v(x)}$ . In general  $\psi$  is strictly positive and only at those points equal to zero, where u is equal to zero. We now conclude from the first equation that  $0 = \nabla(\nabla \psi e^v)$  and therefore  $\psi$  satisfies the equation  $\Delta \psi + \nabla \psi \nabla v = 0$  in  $\Omega \subset \mathbb{R}^2$  with Neumann boundary data at  $\partial\Omega$ . If we restrict ourselves to functions (u, v) that are both finite everywhere we see that  $\psi = ue^{-v}$ . However the equation for  $\psi$  implies that this function cannot attain a critical point in  $\Omega$ , since at such a point the gradient vanishes and  $\Delta \psi$  would be either strictly positive or strictly negative. According to the boundary conditions Hopf's maximum principle [31, Hopf's Lemma, page 330] implies that  $\psi$  is equal to a constant. Thus  $u(x) = \operatorname{const} \cdot e^{v(x)}$ . This however implies that u and v attain their extrema at the same point in  $\Omega$ , since u is a monotonic function of v and as a consequence from the first equation of (9) we see that  $\nabla u - u\nabla v = 0$ , i.e. the population current vanishes everywhere in  $\Omega$ . A result, independent from the reaction terms of the second equation.

However this result contains the assumption that the functions u and v are finite everywhere in  $\Omega$  and therefore  $\psi$  is finite in  $\Omega$ . So if the time dependent equations describe aggregation, such an assumption then has to break down in the points where the aggregation takes place, i.e. in the aggregation centers. From the fact that for the time dependent problem the  $L^1$ -norm of the solutions is uniformly bounded by the  $L^1$ -norm of the initial data we see that the set of points where aggregation takes place has to be a set of measure zero.

In [106] V. Nanjundiah also elaborated the fact that the singularities can only be of  $\delta$ -function type. Therefore he remarked that the trivial solution satisfies the equations (9) pointwise. The mass condition on the solution can only be satisfied if the solution has singularities. Since we have from the previous arguments that  $u = Ke^v$  we get for v the problem  $\Delta v + Ke^v - v = 0$  with homogeneous Neumann boundary data on  $\partial\Omega$ . From this equation one can derive all possible steady state solutions. Furthermore a uniform solution  $v \equiv const =: L$  always exists, where L is defined by the mean value of v over  $\Omega$ .



Figure 1: The connection between the values of the constants K and L.

Figure 1 shows the connection between K and L. We see that possible non-constant steady state solutions can have values of K in the range of the interval  $[0, Le^{-L}]$ . For K = 0 there are two possible values, namely L = 0 and  $L \to \infty$ .

Furthermore V. Nanjundiah showed that all solutions between the one with K = 0 and the uniform solution are unstable if the uniform one itself is unstable by imposing small time dependent fluctuations on an arbitrary solution at time t = 0. In view of (8) the uniform solution is unstable if  $u = v = v_0$  with  $v_0 > 1$  is true.

Nanjundiah's paper was followed by two papers which contain conjectures for the asymptotic behaviour of the solution of the Keller-Segel model for the space dimensions N = 1, N = 2 and  $N \ge 3$ . In [24] S. Childress and J. K. Percus pointed out that the arguments used by V. Nanjundiah are independent of the dimension of space in which aggregation occurs. However they showed that singular behaviour of the solution is in fact a phenomenon that is space dependent. In their paper they restricted themselves to the (as they called it) minimal system given by the equations

(10) 
$$\begin{cases} u_t = \nabla(\nabla u - \chi u \nabla v), & x \in \Omega, \ t > 0\\ v_t = k_c \Delta v - \gamma v + \alpha u, & x \in \Omega, \ t > 0\\ 0 = \partial u / \partial n = \partial v / \partial n, & x \in \partial \Omega, \ t > 0\\ u(0, x) = u_0(x), \ v(0, x) = v_0(x), & x \in \Omega, \end{cases}$$

which (as mentioned before) is due to some simplifying assumptions done by V. Nanjundiah in [106] and is nowadays the most common formulation of the chemotaxis equations. Their studies and their performed asymptotic expansion analysis (see [24, page 236–237]) lead to the following possible time asymptotic behaviour for the solution of system (10):

"In particular, for the special model we have investigated, collapse cannot occur in a one-dimensional space; may or may not in two dimensions, depending upon the cell population; and must, we surmise, in three or more dimensions under a perturbation of sufficiently high symmetry."

Übersichtsartikel	Historischer Artikel	Buchbesprechungen
		Bachbooproonlangon

Here Childress and Percus refer that aggregation proceeds to the formation of  $\delta$ -functions in the cell density as *chemotactic collapse*. Their analysis and conjecture for N = 2 and  $\Omega$  a disk was confirmed by the result in [25], where S. Childress gives an asymptotic expansion describing the imminent collapse of a radially symmetric aggregate of chemotactic cells. However the studies of the stationary problem continued independently from this conjecture and the report on the time independent problem should be closed first until the time asymptotic behaviour of the solution of (10) becomes the main subject of the present considerations in the upcoming sections.

The papers by Childress and Percus were followed by the studies of stationary solutions done by R. Schaaf. In [128] she analyzed solutions of the system

(11) 
$$\begin{cases} 0 = \nabla(k_1(u,v)\nabla u - k_2(u,v)\nabla v), & x \in \Omega, \ t > 0 \\ 0 = k_c\Delta v + g(u,v), & x \in \Omega, \ t > 0 \\ \partial u/\partial n = \partial v/\partial n = 0, & x \in \partial\Omega, \ t > 0 \\ u(0,x) = u_0(x), \ v(0,x) = v_0(x), & x \in \Omega. \end{cases}$$

with general nonlinearities satisfying the conditions

- 1.  $\Omega \subset \mathbb{R}^N$  is a bounded open region with smooth boundary.
- 2.  $k_1, k_2 : \mathbb{R}^+ \times \mathbb{R}^+ \to \mathbb{R}^+$  are twice continuously differentiable and the ODE

$$\frac{d}{ds}r(s) = k_2(r,s)/k_1(r,s)$$

has a unique solution  $r: \mathbb{R}^+ \to \mathbb{R}^+$  for any initial condition  $r(s_0) = r_0$ ,  $s_0, r_0 \in \mathbb{R}^+$ .

3.  $g: \mathbb{R}^+ \times \mathbb{R}^+ \to \mathbb{R}$  is twice continuously differentiable and  $g^{-1}(\{0\}) \neq \emptyset$ .

via bifurcation techniques. Furthermore a criterion for bifurcation of stable nonhomogeneous aggregation patterns is given. In [128] R. Schaaf focused on the properties of stationary solutions of the Keller-Segel model with homogeneous Neumann boundary data in a very general setting. She shows that the stationary problem of the Keller-Segel model in a more general setting than the cases studied by V. Nanjundiah can also be reduced to a parameter-dependent single scalar equation. More precisely she shows the following theorem:

**Theorem 1 (Schaaf)** A pair  $(u, v) \in \{w \in \mathcal{X} \mid w(\overline{\Omega}) \subset \mathbb{R}^+\} \times \{w \in \mathcal{X} \mid w(\overline{\Omega}) \subset \mathbb{R}^+\}$  is a solution of (11) iff, for  $\lambda \in \mathbb{R}^+$ ,

(12)  $u(x) = \varphi(v(x), \lambda)$  for all  $x \in \overline{\Omega}$  and  $k_c \Delta v + g(\varphi(v(x), \lambda), v) = 0$ .

Here the space  $\mathcal{X}$  is defined as  $\{w \in \mathbb{Z} \mid \partial w / \partial n = 0\}$  where  $\mathbb{Z}$  is the space  $C^{2,\beta}(\overline{\Omega}, \mathbb{R})$  with  $0 < \beta < 1$  for N > 1 and  $C^2(\overline{\Omega}, \mathbb{R})$  for N = 1. The function  $\varphi(s, \lambda)$  is given by r(s) with

$$\frac{d}{ds}r(s) = k_2(r,s)/k_1(r,s), \ r(1) = \lambda.$$

Then bifurcation methods are used in [128] to find natural bifurcation points. Furthermore R. Schaaf gives a stability analysis for the constant stationary solutions of the Keller-Segel model. For  $k_1(u, v) = 1$ ,  $k_2(u, v) = \chi u$  and  $g(u, v) = -\gamma v + \alpha u$  the stationary solutions of the Keller-Segel model solve the equation

(13) 
$$k_c \Delta v - \gamma v + \alpha \lambda \exp(\chi v) = 0$$
 in  $\Omega$ 

with homogeneous Neumann boundary data. Of course the question of positive, nontrivial solutions for this equation arises. The existence of nontrivial radially symmetric solutions for this equation has been shown in [11, Proposition 1] under the assumption that  $\gamma > 0$ , but Biler did not consider the nonsymmetric case (The Neumann boundary data implies that there are no solutions provided  $\gamma = 0$ .).

Using variational techniques introduced by M. Struwe and G. Tarantello in [147] J. Wei and G. Wang [153], T. Senba and T. Suzuki [133, 134] and in [63] myself proved for  $\Omega \subset \mathbb{R}^2$  independently the existence of nontrivial solutions of (13) without symmetry assumptions for  $4\pi < \alpha \chi \lambda/k_c$ . The existence of nontrivial solutions of (13) in the case that  $\alpha \chi \lambda/k_c < 4\pi$  follows from arguments that will be mentioned later in the present paper.

The idea of the existence proof is based on the studies of the functional

$$\mathcal{F}_{\alpha\chi\lambda/k_c}(v) := \frac{1}{2} \int_{\Omega} |\nabla v|^2 + \gamma v^2 dx - \frac{\alpha\chi\lambda}{k_c} \log\left(\frac{1}{|\Omega|} \int_{\Omega} e^v dx\right),$$

where  $v \in \mathcal{D} := \{v \in H^1(\Omega) \mid v \text{ has mean value equal to zero over } \Omega\}$ . One easily notices that  $v \equiv 0$  is a strict local minimum for  $\mathcal{F}_{\alpha\chi\lambda/k_c}$  in the case that

$$\gamma > \frac{\alpha \chi \lambda}{|\Omega| k_c} - \mu_1,$$

where  $\mu_1$  is the first (non zero) eigenvalue of the Laplacian with homogeneous Neumann boundary data. Then one recognizes that for a smooth domain  $\Omega$  and  $\alpha \chi \lambda > 4k_c \pi$  there is a sequence  $\{v_{\varepsilon}\}_{\varepsilon>0} \subset \mathcal{D}$  with

$$v_{\varepsilon}(x) = \log\left(\frac{\varepsilon^2}{\left(\varepsilon^2 + \pi |x - x_0|^2\right)^2}\right) - \frac{1}{|\Omega|} \int_{\Omega} \log\left(\frac{\varepsilon^2}{\left(\varepsilon^2 + \pi |x - x_0|^2\right)^2}\right) dx,$$

where  $x_0$  is an arbitrary point on  $\partial \Omega$ , such that

$$\mathcal{F}_{\alpha\chi\lambda/k_c}(v_{\varepsilon}) \to -\infty \text{ and } ||\nabla v_{\varepsilon}||_{L^2(\Omega)} \to \infty \text{ as } \varepsilon \to 0.$$

As a consequence there exists a  $v_0 \in \mathcal{D}$  such that

 $\mathcal{F}_{\alpha\chi\lambda/k_c}(v_0) < 0 \text{ and } ||v_0||_{H^1(\Omega)} \ge 1.$ 

One now defines

 $\mathcal{P} \equiv \{ p : [0,1] \to \mathcal{D} \mid p \text{ is continuous and } p(0) = 0, \ p(1) = v_0 \}$ 

and sets

$$k_{\alpha\chi\lambda/k_c} \equiv \inf_{p\in\mathcal{P}} \max_{t\in[0,1]} \mathcal{F}_{\alpha\chi\lambda/k_c}(p(t))$$

for all  $\alpha \chi \lambda / k_c \ge 4\pi$ . Using the fact that the mapping

$$\alpha \chi \lambda / k_c \mapsto \frac{k_c k_{\alpha \chi \lambda / k_c}}{\alpha \chi \lambda}$$

is monotone decreasing for all  $\alpha \chi \lambda/k_c \ge 4\pi$  we see that it is differentiable for almost every  $\alpha \chi \lambda/k_c \ge 4\pi$ . The rest of the proof then consists of the construction of a Palais-Smale sequence for  $\mathcal{F}_{\alpha \chi \lambda/k_c}$  that contains a subsequence that converges strongly in  $H^1(\Omega)$  to a critical point of  $\mathcal{F}_{\alpha \chi \lambda/k_c}$ . The construction of the Palais-Smale sequence can be done exactly as in the paper by M. Struwe and G. Tarantello [147]. The existence of the nontrivial critical point of the Functional  $\mathcal{F}_{\alpha \chi \lambda/k_c}$  over the set  $\mathcal{D}$  allows us to conclude the existence of a nontrivial solution of equation (13). This can be seen easily. If one introduces the new function

$$w := \chi v - \frac{\chi}{|\Omega|} \int_{\Omega} v \, dx$$

we get from (13) the Euler-Lagrange equation of the minimizing problem  $\inf \mathcal{F}_{\alpha\chi\lambda/k_c}(\nu)$  over the set  $\mathcal{D}$ . Thus the existence of a nontrivial critical point of the functional gives us also the existence of a nontrivial steady state solution of the Keller-Segel model with a linear sensitivity function.

With different methods than those just mentioned Y. Kabeya and W.-M. Ni also proved the existence of positive nontrivial stationary solutions of (13) in [70]. Furthermore they showed the following result:

**Theorem 2 (Kabeya & Ni)** Let  $\Omega \subset \mathbb{R}^2$ . Suppose that  $t = \lambda e^{\chi t}$  has two positive solutions. Then there exists a non constant solution  $v_{\varepsilon}$  of (13) provided

$$arepsilon:=\sqrt{k_c/\gamma}$$

is sufficiently small. Moreover, there exist constants  $C_1 > 0$ ,  $C_2 > 0$ ,  $\delta > 0$ , K > 0 and  $\varepsilon > 0$  such that:

$$\sup_{\Omega} v_{\varepsilon} \leq C_{1}, \inf_{\Omega} v_{\varepsilon} \leq C_{2} e^{-\frac{\delta}{\varepsilon}} and \int_{\Omega} (\varepsilon^{2} |\nabla v_{\varepsilon}|^{2} + \gamma v_{\varepsilon}^{2}) dx \geq K \varepsilon^{2}$$

for any  $0 < \varepsilon < \varepsilon_0$ . Furthermore for sufficiently small  $\varepsilon > 0$  the solution  $v_{\varepsilon}$  has exactly one local maximum point in  $\overline{\Omega}$ , which must lie on the boundary  $\partial\Omega$ .

This theorem is similar to results that have been established for the stationary Keller-Segel model with a logarithmic chemotactical sensitivity. In this case the transformation introduced by R.Schaaf in [128] leads to the problem

(14)  $d\Delta w - w + w^p = 0$ , in  $\Omega$  with  $\partial w / \partial n = 0$ , on  $\partial \Omega$ .

In [83, 108] and [109] the authors prove the existence of stationary solutions of this equation for  $\Omega \subset \mathbb{R}^N$  with  $N \ge 2$  and  $1 if <math>N \ge 3$  and 1 if <math>N = 2. Their results for this problem can be summarized as follows:

**Theorem 3 (Ni & Takagi)** Let  $w_d$  be a least energy solution of (14), i.e. a critical point of

$$J_d(v) := \int_{\Omega} \frac{1}{2} (d|\nabla v|^2 + v^2) - \frac{1}{p+1} v_+^{p+1} dx$$

such that

$$J_d(w_d) = c_d$$
 where  $c_d := \inf_{h \in \Gamma} \max_{0 \le t \le 1} J_d(h(t))$ ,

in which

$$\Gamma := \{ h \in C([0,1]; W^{1,2}(\Omega)) \mid h(0) = 0, h(1) = e \}$$

and  $e \neq 0$  is a nonnegative function in  $W^{1,2}(\Omega)$  with  $J_d(e) = 0$ . Then  $w_d$  has at most one local maximum in  $\overline{\Omega}$  and this is attained in exactly one point which must lie on the boundary, provided that d is sufficiently small. If  $P_d \in \partial \Omega$  is the unique point at which  $\max w_d$  is achieved. Then

 $\lim_{d\to 0} H(P_d) = \max_{P\in\partial\Omega} H(P)$ 

where H(P) denotes the mean curvature of  $\partial \Omega$  at P.

Of course many generalizations of this result have been published in the recent years (see for example the papers by [47] and [123]), but it is not the goal of the present paper to mention all these results. Therefore I leave this to the interested reader and turn now to the time dependent problem. However it is recommended to recall the presented results when looking at the time asymptotic behaviour of the solution in the upcoming section. Recalling the results of the present section will help to understand the results for the time asymptotics of the solution and will help to understand which behaviour one might expect for the solution. Before we now definitively turn to the time dependent problem let us explain some terms used in the present section. We have seen that there are different effects that one can expect. In some cases we spoke of aggregation and in other cases of a special form of aggregation namely the formation of  $\delta$ -singularities. This was sometimes called chemotactical collapse. Before we turn to the time dependent model we therefore now introduce three important effects in the mathematical language.

**Definition 1** Let (u(t, x), v(t, x)) be a solution of (4) for the corresponding initial data  $(u_0(x), v_0(x))$ . We say that the model describes **aggregation**, if

 $\liminf_{t \to \infty} ||u(t, \cdot)||_{L^{\infty}(\Omega)} > ||u_0(\cdot)||_{L^{\infty}(\Omega)}$ 

and  $||u(t, \cdot)||_{L^{\infty}(\Omega)} < \text{konst for all } t$ . The solution **blows up** resp. is a **blow-up solution** if  $||u(t, \cdot)||_{L^{\infty}(\Omega)}$  or  $||v(t, \cdot)||_{L^{\infty}(\Omega)}$  becomes unbounded in either finite or infinite time, i.e. there exists a time  $T_{\max}$  with  $0 < T_{\max} \leq \infty$  such that

$$\limsup_{t \to T_{\max}} ||u(t, \cdot)||_{L^{\infty}(\Omega)} = \infty \ or \ \limsup_{t \to T_{\max}} ||v(t, \cdot)||_{L^{\infty}(\Omega)} = \infty.$$

We will speak of chemotactical collapse if  $\limsup ||u(t, \cdot)||_{L^{\infty}(\Omega)} < ||u_0(\cdot)||_{L^{\infty}(\Omega)}$ .

$$t \rightarrow \infty$$

Übersichtsartikel	Historischer Artike
Obcidicitiourinci	

Remark that this definition is almost identical to that given in [116] beside the difference in the allowed blow-up time for a blow-up solution. Furthermore remark that the three cases do not exclude themselves, i.e. more than one case can happen for the same solution. At the end of this section let us summarize the mentioned results in the following table:

Table 1: Collection of results for the stationary Keller-Segel models.

Observation	References
For model (4) the uniform distribution $(u_0, v_0)$ becomes unstable if $\frac{k_2(u_0, v_0)v_0}{k_1(u_0, v_0)u_0} + \frac{u_0 f'(v_0)}{k_3(v_0) + v_0 k'_3(v_0)} > 1.$	[71]
All solutions between the one with $K = 0$ (as defined in this section) and the uniform solution are unstable if the uniform one itself is. unstable.	[106]
The stationary problem of the Keller-Segel model can be reduced to the parameter-dependent single scalar equation (12).	[106, 128]
There exist non-constant stationary solutions of the Keller-Segel model for example for a linear and for a logarithmic chemotactical sensitivity function.	[11, 63, 70, 83] [108, 109, 133] [134] and [153]

## 4 The time dependent problem: The case of a linear chemotactical sensitivity

The conjectures and observations by V. Nanjundiah, S. Childress and J. K. Percus have been the initiating motivations for a large number of researchers to study the time asymptotic behaviour of the solution of the system (10). There is still an avalanche of publications running and I am pretty sure that by the date of publication of the present paper the number will have increased once again. Parallel to the results of this section various papers were published in which versions of the Keller-Segel model with a different chemotactic sensitivity function were studied. I will mention these results in an upcoming section. Thus we will only focus on results for (10) in this section resp. the upcoming subsections.

## 4.1 Early results on the time asymptotics and conjectures

The first step in the analysis of the conjectures by Childress and Percus has been done by W. Jäger and S. Luckhaus in 1992. In [69] they introduced the transformation

$$U(t,x) := \frac{|\Omega|u(t,x)}{\int\limits_{\Omega} u(t,x)dx} \text{ and } V(t,x) := v(t,x) - \frac{1}{|\Omega|} \int\limits_{\Omega} v(t,x)dx$$

which leads to the system:

(15) 
$$\begin{cases} U_t = \nabla(\nabla U - \chi U \nabla V), & x \in \Omega, \ t > 0\\ \frac{1}{k_c}(V_t + \gamma V) = \Delta V + \frac{\alpha}{k_c}(U - 1), & x \in \Omega, \ t > 0\\ \frac{\partial U}{\partial n} = \frac{\partial V}{\partial n} = 0, & x \in \partial\Omega, \ t > 0\\ U(0, x) = U_0(x), \ V(0, x) = V_0(x), & x \in \Omega. \end{cases}$$

Jäger and Luckhaus then assume that  $\alpha = k_c\beta$ , the constants  $\chi$ ,  $k_c$ ,  $\alpha$  are of the order  $\frac{1}{\varepsilon}$  with  $\varepsilon$  small and  $\gamma$  and  $\beta$  are of order 1. Thus they get for small  $\varepsilon$  resp. for  $\varepsilon \to 0$  the system

(16) 
$$\begin{cases} U_t = \nabla(\nabla U - \chi U \nabla V), & x \in \Omega, \ t > 0\\ 0 = \Delta V + \frac{\alpha}{k_c}(U-1), & x \in \Omega, \ t > 0\\ \partial U/\partial n = \partial V/\partial n = 0, & x \in \partial \Omega, \ t > 0\\ U(0, x) = U_0(x) & x \in \Omega. \end{cases}$$

Their result in space dimension N = 2 for system (16) is summarized as follows:

**Theorem 4 (Jäger & Luckhaus)** Let  $\Omega$  be a bounded open set in  $\mathbb{R}^2$ ,  $\partial\Omega$  is a  $C^1$ -boundary,  $U_0(x)$  is  $C^1$  and satisfies the boundary condition.

- 1. There exists a critical number  $c(\Omega)$  such that  $\beta \chi \overline{U_0(x)} < c(\Omega)$  implies that there exists a unique, smooth positive solution to (16) for all time.
- 2. Let  $\Omega$  be a disk. There exists a positive number  $c^*$  with the following property: If  $\beta \chi \overline{U_0(x)} > c^*$  then radially symmetric positive initial values can be constructed such that explosion of U(t, x) happens in the center of the disk in finite time.

Here the notation  $\overline{U_0(x)}$  is used for the mean value of  $U_0(x)$  over the domain  $\Omega$ . More precisely Jäger and Luckhaus show the following Proposition which implies *1*. of the previous Theorem 4.

**Proposition 1 (Jäger & Luckhaus)** Let  $\Omega$  be a domain satisfying the smoothness assumptions from the Theorem above. Let U(t, x) be a smooth positive solution to (16) and  $t^*$  the maximal time of existence,  $0 < t^* \leq \infty$ . There exists a positive number  $c_1(\Omega)$  such that  $t^* < \infty$  implies

$$\lim_{k\to\infty}\overline{\lim_{t\to t^*}}\beta\chi\overline{U_0(x)}\int_{\Omega}(U(t,x)-k)_+dx\geq c_1(\Omega).$$

Proposition 1 is shown by multiplying the first equation of (16) with  $\varphi = (U - k)_+^{m-1}$ where  $k \ge 0$  and m > 1. Then the second equation of (16) allows to estimate the term

$$-\int_{\Omega} U\nabla V\nabla (U-k)_{+}^{m-1} dx = -\int_{\Omega} \nabla V\nabla \left(\frac{m-1}{m}(U-k)_{+}^{m} + k(U-k)_{+}^{m-1}\right) dx$$

from above by

$$c(k, m) \int_{\Omega} (U-k)_+^m dx.$$

Übersichtsartikel	Historischer Artikel	Buchbesprechungen
-------------------	----------------------	-------------------

If the statement of Proposition 1 did not hold this estimate would allow us to find the following inequality

$$\frac{d}{dt} \int_{\Omega} (U-k)_+^m dx \le c_2(\Omega,k) + c_3(k) \int_{\Omega} (U-k)_+^m dx \text{ for all } t \ge t_1$$

which would give us a bound for the  $L^m$ -norms of U(t, x) and the global existence would follow by standard regularity arguments for solutions of elliptic and parabolic equations.

For the proof of the blow-up statement 2. of Theorem 4, W. Jäger and S. Luckhaus studied the function

$$M(t,\rho) := \int_{0}^{\sqrt{\rho}} (U(t,r) - 1)r \, dr \text{ for } r = |x|, \ 0 \le \rho \le R.$$

Using the equations of (16) they found that  $M(t, \rho)$  has to solve the following initial boundary value problem:

$$\frac{\partial}{\partial t}M = 4\rho \frac{\partial^2}{\partial \rho^2}M + \beta \chi \overline{U_0(x)} \frac{\partial}{\partial \rho}M + \beta \chi \overline{U_0(x)}M,$$

with

$$M(0,\rho) = \int_{0}^{\sqrt{p}} (U_0(r) - 1)r \, dr \text{ and } M(t,0) = M(t,R) = 0.$$

Constructing a subsolution  $W(t, \rho)$  for this problem such that  $W(t, \rho) \le M(t, \rho)$  for all  $t, \rho$  and

$$\lim_{t \to T_{finite}} \sup_{\rho < \epsilon} W(t, \rho) \ge \omega > 0$$

10

for each  $\epsilon > 0$  they proved that the solution has to blow up at time  $T_{finite}$  in the center of the disk. Furthermore Jäger and Luckhaus asked for more information about the blow-up behaviour of the solution of (16). In a remark [69, page 820] they formulated the following questions:

"It would be interesting to know more about the set of explosion points at  $t^*$ . The solution may globally exist as weak solutions. The development of singularities after a finite time  $t^*$  is another important topic to be studied."

Even though it was not the next paper in the chronology I now mention the results from M. A. Herrero and J. J. L. Velázquez [50] from 1996 and M. A. Herrero, E. Medina and J. J. L. Velázquez [53, 54] from 1997 and 1998 since they studied system (16) in those papers. In [50, 55] they focused on the possible formation of  $\delta$ -function singularities in finite time in space dimension N = 2. Using asymptotic expansion methods in [50] their result was the following: **Theorem 5 (Herrero & Velázquez)** Let R > 0, and let  $\Omega_R = \{x \in \mathbb{R}^2 : |x| < R\}$ . Then there exist radial solutions of (16) defined in an interval (0, T) with T > 0, and such that:

(17) 
$$U(t,r) \rightarrow \frac{8\pi k_c}{\chi \alpha} \delta(0) + \psi(r) \text{ as } t \rightarrow T,$$

in the sense of measures, where  $\delta(0)$  is the Dirac measure centered at r = 0, and:

(18) 
$$\psi(r) = \frac{C}{r^2} e^{-2|\log(r)|^{1/2}} (2|\log(r)|)^{\frac{1}{2\sqrt{2}|\log(r)|^{1/2}} - \frac{1}{2}} (1 + o(1))$$

as  $r \to 0$ , where C is a positive constant depending on  $\chi$ . At t = T, the profile near r = 0 is given by:

(19) 
$$U(t,r) = \frac{8\pi k_c}{\chi \alpha} \delta(0) + \psi(r); \ \psi(r) \ as \ in \ (18).$$

Moreover, if we set  $S(t) = (T - t)(\sup_{\Omega} U(t, r)) \equiv (T - t)U(0, t)$ , one has that  $\lim_{t \to T} S(t) = \infty$ . More precisely, there holds:

$$S(t) = C_1(T-t)^{-1} |\log(T-t)|^{1-\frac{1}{\sqrt{|\log(T-t)|}}}$$
 as  $t \to T$ , for some  $C_1 > 0$ 

So they found solutions that form in finite time a  $\delta$ -singularity in the center of the disk in  $\mathbb{R}^2$ . Furthermore they investigated a result for the whole space in the three dimensional case in [53, 54] and [55]. There they studied self-similar solutions and could formulate the following statements.

#### Theorem 6 (Herrero, Medina & Velázquez)

1. Consider (16) in space dimension N = 3 with  $\Omega = \mathbb{R}^3$ . Then, for any T > 0 and any constant C > 0, there exists a radial solution (U(t,r), V(t,r)) of (16) that is smooth for all times 0 < t < T, blows up at r = 0 and t = T, and is such that:

$$\int_{|x|\leq r} U(T,s) \, ds \to C.$$

2. Consider (16) in space dimension N = 3 with  $\Omega = \mathbb{R}^3$ . For any T > 0 there exists a sequence  $\{\delta_n\}_{n \in \mathbb{N}}$  with  $\lim_{n \to \infty} \delta_n = 0$ , and a sequence of radial solutions  $(U_n(t,r), V_n(t,r))$ , that blows up at r = 0 and t = T, and are such that  $U_n(t,r)$  is self-similar, and

$$U_n(t,r) \sim \left(\frac{8\pi k_c}{\chi lpha} + \delta_n\right) (4\pi r^2)^{-1} \text{ as } r \to 0.$$

For this solution

$$N(t,r) := \int\limits_{|x| \le r} U(T,s) \; ds o 0 \; ext{as} \; r o 0$$

Übersichtsartikel	Historischer Artikel	Buchbesprechungen
		_ a chine copie chinan gen

3. No radial, self-similar solution of (16) exists such that  $N(t,r) < \infty$  as  $r \to 0$  when N = 2, resp. when  $\Omega = \mathbb{R}^2$ .

Thus one slowly got more and more insights for the Keller-Segel model but at this point several questions still remained open. To name a few beside the questions raised in [69] we list the following.

- 1. What happens if one drops the assumption of radial symmetry of the solution and how does in the case of blow-up the blow-up profile of the solutions look like?
- 2. Is it possible to prove blow up results also for the full system (10)?
- 3. Can one give the precise value of the threshold which decides whether the solution might blow up or not?

As in the section before I summarize the results of this section in a table, too.

Table 2: Possible time asymptotical behaviour of the solutions of the simplified model (16).

Dimension	Observation	References
N = 2	There exists a critical value $c(\Omega)$ such that a unique, smooth positive solution to (16) exists globally in time if $\beta \chi \overline{U_0(x)} < c(\Omega)$ .	[69]
	Let $\Omega$ be a disk. Then there exists a positive number $c^*$ such that there exists radially symmetric positive initial data with the following property: If $\beta \chi U_0(x) > c^*$ then radially symmetric positive initial values can be constructed such that explosion of $U(t, x)$ happens in the center of the disk in finite time.	
	There exists radially initial data such that the solution of (16) forms in the center of a disk $\Omega$ a $\delta$ -function singularity described in Theorem 5 in finite time.	[50, 55]
	When $\Omega = \mathbb{R}^2$ , then no radial, self-similar solutions of (16) exist such that $\int_{ x  \le r} U(T, s)  ds < \infty  \text{as}r \to 0.$	[53, 55]
<i>N</i> = 3	Let $\Omega = \mathbb{R}^3$ . Then there exists, for any $T > 0$ and any constant $C > 0$ , a radial solution $(U(t, r), V(t, r))$ of (16) that is smooth for all times $0 < t < T$ , blows up at $r = 0$ and $t = T$ , and is such that: $\int_{ x  \le r} U(T, s) ds \to C$ .	[53, 54, 55]
	For any $T > 0$ there exists a sequence $\{\delta_n\}_{n \in \mathbb{N}}$ with $\lim_{n \to \infty} \delta_n = 0$ , and a sequence of radial solutions $(U_n(t,r), V_n(t,r))$ , that blows up at $r = 0$ and $t = T$ , and are such that $u_n(t,r)$ is self-similar, and $U_n(t,r) \sim \left(\frac{8\pi k_c}{\chi \alpha} + \delta_n\right) (4\pi r^2)^{-1}$ as $r \to 0$ . For this solution $\int_{ x  \leq r} U(T,s) ds \to 0$ as $r \to 0$ .	

#### 4.2 Progress and further questions

After W. Jäger's and S. Luckhaus' paper in 1992 the next step was performed 1995 by T. Nagai in [92]. In his article "*Blow-up of radially symmetric solutions to a chemotaxis system*" [92] he proved the following result for the simplified system

(20) 
$$\begin{cases} u_t = \nabla(\nabla u - \chi u \nabla v), & x \in \Omega, \ t > 0\\ 0 = \Delta v - \gamma v + \beta u, & x \in \Omega, \ t > 0\\ \partial u / \partial n = \partial v / \partial n = 0, & x \in \partial \Omega, \ t > 0\\ u(0, x) = u_0(x), & x \in \Omega. \end{cases}$$

#### Theorem 7 (Nagai)

*1.* Suppose that N = 1, or N = 2 and

$$\beta \chi \int\limits_{B(0,R)} u_0(x) dx < 4\omega_N$$

with radially symmetric 
$$u_0(x)$$
. Then  $T_{\max} = \infty$  and  

$$\sup_{t \ge 0} \left\{ ||u(t, \cdot)||_{L^{\infty}(B(0,R))} + ||v(t, \cdot)||_{L^{\infty}(B(0,R))} \right\} < \infty.$$

*2.* Let  $N \ge 2$  and  $u_0$  be radially symmetric. If

$$\begin{split} 0 &> 2N(N-1) \left( \frac{1}{\omega_N} \int_{\Omega} u_0(x) dx \right)^{2/N} \left( \frac{1}{\omega_N} \int_{B(0,R)} u_0(x) |x|^N dx \right)^{(N-2)/N} \\ &- \frac{N}{2} \beta \chi \left( \frac{1}{\omega_n} \int_{\Omega} u_0(x) dx \right)^2 \\ &+ \beta \chi N R^{-N} \left( \frac{1}{\omega_n} \int_{\Omega} u_0(x) dx \right) \left( \frac{1}{\omega_N} \int_{B(0,R)} u_0(x) |x|^N dx \right) \\ &+ \beta \chi \gamma \begin{cases} \frac{1}{e} \left( \frac{1}{\omega_N} \int_{\Omega} u_0(x) dx \right)^{3/2} \left( \frac{1}{\omega_N} \int_{B(0,R)} u_0(x) |x|^N dx \right)^{1/2}, \\ if \ N = 2 \\ \frac{1}{2(N-2)} \left( \frac{1}{\omega_N} \int_{\Omega} u_0(x) dx \right)^{(2N-2)/N} \left( \frac{1}{\omega_N} \int_{B(0,R)} u_0(x) |x|^N dx \right)^{2/N}, \\ if \ N \ge 3 \end{split}$$

where  $\omega_N$  denotes the area of the unit sphere  $S^{N-1}$  in  $\mathbb{R}^N$ , then  $T_{\max} < \infty$  and  $\limsup_{t \to T_{\max}} ||u(t, \cdot)||_{L^{\infty}(B(0,R))} = \infty.$  Furthermore the radially symmetric solution (u(t,r), v(t,r)) of (20) satisfies

$$u(t,r) + v(t,r) \le K(n)$$
 for  $\frac{1}{n} \le r \le R$  and  $0 \le t < T_{\max}$ 

where K(n) denotes a generic positive constant depending on  $n \in \mathbb{N}$  such that

 $K(n) \to \infty$  as  $n \to \infty$ .

Thus the blow-up can only occur at the point r = 0.

While the first statement is easy to check the second is based on some subtle estimates of the expression

$$M_N(t) := \frac{1}{\omega_N} \int\limits_{B(0,R)} u(t,x) |x|^N dx.$$

The global existence proof of solutions of (20) in one space dimension performed in [92] illustrates in a nice way how one tries to show the existence of the solution global in time in higher space dimensions. Therefore I first want to demonstrate this proof here. If one integrates for N = 1 the second equation of (20) on (-R, x) one gets

$$v_x(t,x) = \gamma \int_{-R}^{x} v(t,y) dy - \beta \int_{-R}^{x} u(t,y) dy.$$

Thus we see that

$$|v_x(t,x)| \leq \beta \int_{-R}^{R} u_0(x) dx \text{ on } \Omega \times (0,T_{\max}).$$

For  $x \in \Omega = (-R, R)$  we now obtain

$$2Rv(t,x) = \int_{-R}^{R} v(t,y) dy + \int_{-R}^{R} \int_{y}^{x} v_{x}(t,z) dz \, dy$$

and therefore

$$0 \leq v(t,x) \leq \frac{\beta}{2R} \left(\frac{1}{\gamma} + 4R^2\right) \int_{\Omega} u_0(x) dx.$$

Thus

 $||v(t, \cdot)||_{L^{\infty}(\Omega)} \leq const$  and  $||v_x(t, \cdot)||_{L^{\infty}(\Omega)} \leq const$  for all  $0 < t < T_{\max}$ .

Multiplying now the first equation of (20) with  $u^p$  for  $p \ge 1$  and integrating the equation over  $\Omega$  yields

$$\frac{1}{p+1}\frac{d}{dt}\int_{\Omega}u^{p+1}dx = \frac{-2p}{p+1}\int_{\Omega}|\nabla u^{(p+1)/2}|^2dx + \frac{p(p+1)}{2}\cdot const\cdot\int_{\Omega}u^{p+1}dx.$$

Now the bound of the  $L^{\infty}$ -norm of the solution can be obtained by application of N. D. Alikakos' version of the Moser iteration introduced in [2]. One therefore sees that the question whether the solution exists globally in time or not depends crucially on a uniform bound for the  $L^{\infty}$ -norm of the gradient of v. This simplified version of the Keller-Segel model has been extensively studied by Nagai and his coauthors. Once again we cannot follow the chronology since the different versions of the Keller-Segel model have been studied parallel. Thus I concentrate on the results on system (20) in this subsection and turn to the results for the full Keller-Segel model (10) resp. (15) later on. The simplified versions allow to decouple the system. Therefore techniques are available in these cases which are not at hand for the full parabolic version. For the simplified version (20) recent results from Nagai, Senba, Suzuki et al. give more information about the blow-up profile of the solution and the non symmetric blow up. However their proofs are very technical and desire fine estimates that are difficult to demonstrate in a simple way. Thus I restrict myself to present their results in Table 3 and 4.

Table 3: Possible time asymptotical behaviour of the solutions of the simplified model (20) with  $\gamma>0.$ 

Dimension	Observation	References
N = 1	The solution of the Keller-Segel model exists globally in time and is uniformly bounded for all $t \ge 0$ .	[92]
N = 2	If $\beta \chi \int_{\Omega} u_0(x) dx < 4\pi$ then the classical solution of	[92, 93, 94]
	the Keller-Segel model exists globally in time and is uniformly bounded for all $t \ge 0$ . If $\Omega$ is a disk and $u_0$ is radially symmetric or satisfies $u(x) = u(-x)$ in $\Omega$ , then this statement holds if $\beta \chi \int_{\Omega} u_0(x) dx < 8\pi$ .	[95, 98, 101] and [130]
	Let $x_0 \in \Omega$ . If $\beta \chi \int_{\Omega} u_0(x) dx > 8\pi$ and if $\int_{\Omega} u_0(x)  x - x_0 ^2 dx$ is sufficiently small, then the corresponding solution of (20)	[102]
	and (16) blows up in finite time.	
	Assume that $\partial\Omega$ has a line segment $\mathcal{L}_0$ , and that $\Omega$ lies on one side of a line $\mathcal{L}$ containing $\mathcal{L}_0$ . If furthermore $\beta\chi\int_{\Omega}u_0(x)dx > 4\pi$ and if $\int_{\Omega}u_0(x) x-x_0 ^2dx$	[102]
	is sufficiently small for a point $x_0 \in \mathcal{L}_0$ that is not an end-point of $\mathcal{L}_0$ , then the corresponding solution of (20) and (16) blows up in finite time.	
	If $\Omega$ is a disk, $u_0$ is radially symmetric and if $\beta \chi \int_{\Omega} u_0(x) dx > 4\omega_2$ , then there exists a constant $C_1$	[92]
	depending on $\frac{1}{\omega_2} \int_{\Omega} u_0(x) dx$ such that if	
	$0 < \frac{1}{\omega_2} \int_{\Omega} u_0(x)  x ^2 dx < C_1$ then <i>u</i> blows up in finite time.	

Übersichtsartikel	Historischer Artikel	Buchbesprechungen
-------------------	----------------------	-------------------

Dimension	Observation	References
N = 2	If $\beta \chi \int_{\Omega} u_0(x) dx < 8\pi$ and $T_{\max} < \infty$ then there exists a point in $x_0 \in \partial \Omega$ such that $\limsup_{t \to T_{\max} \Omega \cap B(x_0,\epsilon)} \int u(t,x) dx \ge 2\pi/a_*\beta$ for any $\epsilon > 0$ , where $a_*$ is a root of $a_* - \chi/2 -   u_0  _{L^1(\Omega)}\beta a_*/16\pi = 0$ such that $a_* < \chi$ .	[96, 97]
	Suppose $T_{\max} < \infty$ . Then there exist for any isolated blow-up point $x_0$ two positive constants $\delta, m \ge m_*$ and a non-negative function $f \in L^1(\overline{B(x_0, \delta) \cap \Omega}) \cap C(\overline{B(x_0, \delta) \cap \Omega} \setminus \{x_0\})$ such that $u(t, \cdot)$ converges weakly in the Banach space of all Radon measures on $\overline{B(x_0, \delta) \cap \Omega}$ to $m\delta_{x_0} + f$ as $t \to T_{\max}$ , where $m_*$ is equal to $4\pi/\beta\chi$ if $x_0 \in \partial\Omega$ and equal to $8\pi/\beta\chi$ if $x_0 \in \Omega$ .	[96, 97, 131] and [136]
	Suppose $\Omega = \mathbb{R}^2$ and let $x_0 \in \mathbb{R}^2$ . If $\beta \chi \int_{\mathbb{R}^2} u_0(x) dx > 8\pi$ and if $\int_{\mathbb{R}^2} u_0(x)  x - x_0 ^2 dx$ is sufficiently small, then the corresponding solution of (20) blows up in finite time.	[99]
$N \ge 3$	If $\Omega$ is a sphere and $u_0$ is radially symmetric, then there exists a constant $C_1$ depending on $\frac{1}{\omega_N} \int_{\Omega} u_0(x) dx$ such that if $0 < \frac{1}{\omega_N} \int_{\Omega} u_0(x)  x ^N dx < C_1$ then u blows up in finite time.	[92]
	Suppose $\Omega = \mathbb{R}^N$ and let $x_0 \in \mathbb{R}^N$ . If $\int_{\mathbb{R}^N} u_0(x) x-x_0 ^N dx$ is sufficiently small, then the corresponding solution of (20) blows up in finite time.	[99]

Table 4: Possible time asymptotical behaviour of the solutions of the simplified model (20) with  $\gamma > 0$ .

Of course one can now draw conclusions on the possible number of blow-up points. However I will mention these conclusions a little bit later. Thus at this point let us turn again to a different line of research.

## 4.3 Analysis of the system (15)

Similar to their result for the simplified system (10) M. A. Herrero and J. J. L. Velázquez achieved a very important contribution on the blow-up profile of the solution of the full parabolic systems (10) and the system (15) with  $\gamma = 0$  in their papers [51] and [52]. Using once again asymptotic expansion theory they were able to describe the blowup profile of system (10) and proved therefore the possibility of a  $\delta$ -function formation in finite time for radially symmetric solutions as it was conjectured by Nanjundiah [106] and Childress and Percus [24]. Their main result for system (10) is summarized as follows: **Theorem 8 (Herrero & Velázquez)** Let R > 0, and let  $\Omega_R = \{x \in \mathbb{R}^2 : |x| < R\}$ . Then there exist radial solutions of (10) defined in an interval (0, T) with T > 0, and such that:

(21) 
$$u(t,r) \to \frac{8\pi k_c}{\chi \alpha} \delta(0) + \psi(r) \text{ as } t \to T,$$

in the sense of measures, where  $\delta(0)$  is the Dirac measure centered at r = 0, and:

(22) 
$$\psi(r) = \frac{C}{r^2} e^{-2|\log(r)|^{1/2}} (1 + o(1))$$

as  $r \to 0$ , where C is a positive constant depending on  $\chi$ . At t = T, the profile near r = 0 is given by:

(23) 
$$u(t,r) = \frac{8\pi k_c}{\chi \alpha} \delta(0) + \psi(r); \ \psi(r) \ as \ in \ (22).$$

Moreover, if we set  $S(t) = (T - t)(\sup_{\Omega} u(t, r)) \equiv (T - t)u(0, t)$ , one has that  $\lim_{t \to T} S(t) = \infty$ . More precisely, there holds:

(24) 
$$S(t) = C_1(T-t)^{-1} e^{\sqrt{2|\log(T-t)|}}$$
 as  $t \to T$ , for some  $C_1 > 0$ .

The studies of the asymptotic behaviour of the solution in the non-symmetric case began with the results of [11, 44, 94] and [158]. In [11, 44, 94] the authors introduce independently from each other a Lyapunov functional for system (10) resp. (15) which became an important tool in the then following studies of the time asymptotic behaviour of the solution of system (10) resp. (15). This Lyapunov function is given by

(25) 
$$F(u(t), v(t)) := \int_{\Omega} \frac{k_c}{2\alpha\chi} |\nabla v(t)|^2 + \frac{\gamma}{2\alpha\chi} v^2(t) + u(t)\log(u(t)) - u(t)v(t)dx.$$

Using a Moser-Trudinger type inequality originally formulated by Chang and Yang in [23] the analysis of this functional shows the following:

1. The functional F(u, v) is bounded from below, if

$$\alpha \chi \int\limits_{\Omega} u_0(x) dx \le 4\pi k_c$$

2. The functional F(u, v) is no longer bounded from below, if

$$\alpha\chi\int\limits_{\Omega}u_0(x)dx>4\pi k_c.$$

3. For radially symmetric functions the functional F(u, v) is bounded from below, if

$$\alpha \chi \int\limits_{\Omega} u_0(x) dx \le 8\pi k_c$$

and is no longer bounded from below, if

$$\alpha\chi\int\limits_{\Omega}u_0(x)dx>8\pi k_c.$$

Now two different lines of research became recognizable. One considered system (10) and used more PDE based methods to prove global existence and finite time blow-up results for this system and the other was concerned with system (15) and used methods more related to the calculus of variations. Once again one has to follow these two lines separately to get a clear picture of the achieved results. Let us first have a closer look at the results for (10)

#### 4.3.1 Results for system (10)

Since the question of the well-posedness of a negative cross-diffusion system is not trivial I first turn to the results on the local existence of a solution and possible regularity results. Here one should basically mention A. Yagi [158] and T. Nagai, T. Senba and K. Yoshida [94] whose results can be summarized as follows:

**Theorem 9** Let  $\Omega$  be a bounded, smooth domain in  $\mathbb{R}^2$ . Assume  $u_0, v_0 \in H^{1+\epsilon_0}(\Omega)$  for some  $0 < \epsilon_0 \le 1$  and  $u_0(x) \ge 0$ ,  $v_0(x) \ge 0$  on  $\Omega$ . Let  $T_{\max}$  be the maximal existence time of (u(t), v(t)).

1. (Yagi) System (10) has a non-negative solution (u, v) satisfying  $u, v \in C([0, T_{\max}) : H^{1+\epsilon_1}(\Omega)) \cap C^1((0, T_{\max}) : L^2(\Omega)) \cap C((0, T_{\max}) : H^2(\Omega))$ for any  $0 < \epsilon_1 < \min\{\epsilon_0, 1/2\}$ . Moreover (u, v) has further regularity properties:  $u \in C^1((0, T_{\max}) : H^1(\Omega)), v \in C^{\frac{1}{4}}((0, T_{\max}) : H^3(\Omega)) \cap C^{\frac{5}{4}}((0, T_{\max}) : H^1(\Omega)).$ 

2. (Yagi) If 
$$T_{\max} < \infty$$
, then

$$\begin{split} \lim_{t \to T_{\max}} (||u(t, \cdot)||_{H^{1+\epsilon_0}(\Omega)} + ||v(t, \cdot)||_{H^{1+\epsilon_0}(\Omega)}) &= \infty, \\ \lim_{t \to T_{\max}} \sup_{t \to T_{\max}} ||u(t, \cdot)||_{L^p(\Omega)} &= \infty \text{ for any } 1$$

3. (Nagai, Senba & Yoshida) If

$$\int\limits_{\Omega} u_0(x) dx < \frac{4\Theta k_c}{\alpha \chi}$$

where  $\Theta = 8\pi$  for  $\Omega = \{x \in \mathbb{R}^2 : |x|^2 < R\}$  and  $(u_0, v_0)$  is radial in x and  $\Theta = 4\pi$  otherwise, then the solution (u, v) of (10) exists globally in time and

$$\sup_{t>0}\{||u(t,\cdot)||_{L^{\infty}(\Omega)}+||v(t,\cdot)||_{L^{\infty}(\Omega)}\}<\infty.$$

The local existence and regularity results summarized in Theorem 9 above have been achieved by using semi-group theory. A. Yagi also proved similar local existence results for more general forms of the system (4) in [158] and I will turn to these results later.

The bound of the  $L^{\infty}$ -norm of the solution can once more be achieved by application of N. D. Alikakos' version of the Moser iteration introduced in [2]. Once again the basic and most important step is to find a uniform  $L^{\infty}$ -norm estimate of  $\nabla v(t, \cdot)$  for all  $t \geq 0$ . Nagai, Senba and Yoshida succeeded in finding such a bound in the case where the functional F(u, v) is bounded from below. A. Yagi studied in [158] which norms of the solution have to blow-up if the solution exists only for a finite maximal time of existence  $T_{finite}$ . However beside the results of Herrero and Velázquez in [51, 52] there are no results, that show the existence of initial data such that the corresponding solution of (10) has to blow up in finite time. However there are results under the assumption that there is a solution which blows up in finite time. Let us therefore now turn to those results, that studies the blow-up profile and behaviour of such a solution.

Under the **main assumption** that there is a solution of the Keller-Segel model that blows up in finite time  $T_{finite}$  such that

(26) 
$$\inf_{\substack{0 \le t < T_{finite}}} F(u(t), v(t)) > 0 \text{ or}$$
$$\lim_{t \to T_{finite}} F(u(t), v(t)) = -\infty$$

Nagai, Senba and Suzuki proved in [100, 102] the following results.

**Theorem 10 (Nagai, Senba & Suzuki)** Let  $\Omega \subset \mathbb{R}^2$  be a bounded domain with smooth boundary  $\partial\Omega$ . Furthermore let  $\mathcal{B}$  denote the set of all those points  $x_0$  in  $\overline{\Omega}$  such that there is a sequence  $\{x_k\}_{k\in\mathbb{N}}\subset\overline{\Omega}$  and a sequence  $\{t_k\}_{k\in\mathbb{N}}\subset[0, T_{finite})$  with  $u(t_k, x_k) \to \infty, t_k \to T_{finite}$  and  $x_k \to x_0$  as  $k \to \infty$ .  $\mathcal{B}_I \subset \mathcal{B}$  denotes the set of all isolated blow-up points, i.e  $x_0 \in \mathcal{B}_I$ , iff there exists a R > 0 such that

 $\sup_{0 \leq t < T_{finite}} ||u(t, \cdot)||_{L^{\infty}((B(x_0, R) \setminus B(x_0, r)) \cap \Omega)} < \infty$ 

for any  $r \in (0, R)$  with  $B(x_0, R) := \{x \in \mathbb{R}^2 \mid |x_0 - x| < R\}$ . Then the following statements hold:

1. Given  $x_0 \in \mathcal{B}_I$ , we have 0 < R << 1,  $m \ge m^*$ , and

$$f \in L^1(B(x_0, P) \cap \Omega) \cap C(\overline{B(x_0, R) \cap \Omega} \setminus \{x_0\})$$

satisfying  $f \ge 0$  and  $u(t, \cdot)dx$  converges weakly to  $m\delta_{x_0}(dx) + f dx$  as  $t \to T_{finite}$  in the set of Radon measures on  $\overline{B(x_0, R) \cap \Omega}$ , where

$$m^* := \begin{cases} 8\pi, & x_0 \in \Omega\\ 4\pi, & x_0 \in \partial\Omega. \end{cases}$$

- 2. If (26) occurs, then  $\mathcal{B} = \mathcal{B}_I$ .
- 3. If (10) is radially symmetric and  $T_{\text{max}} < \infty$  then  $\mathcal{B} = \{0\}$ .

These results imply that in the case of a finite time blow-up of the solution the set of isolated blow-up points has finite cardinality and that

$$1 < 2 \times \sharp(\mathcal{B}_I \cap \Omega) + \sharp(\mathcal{B}_I \cap \partial \Omega) \le \frac{\alpha \chi ||u_0||_{L^1(\Omega)}}{4\pi k_c}.$$

Übersichtsartikel	Historischer Artikel	Buchbesprechungen
-------------------	----------------------	-------------------

However a better lower bound of the quantity is of interest in the non radially symmetric case with  $\alpha \chi ||u_0||_{L^1(\Omega)} > 8\pi k_c$ . Where does the blow-up occur? Is there only one blow-up point in the interior of  $\Omega$  or are there two blow-up points at the boundary  $\partial \Omega$  in this case?

Beside the previous results Senba and Suzuki established in [135] the following results using rearrangement and symmetrization arguments. These results are similar to those achieved independently and by other methods in [63] and [64] for system (15).

**Theorem 11 (Senba & Suzuki)** Let  $\Omega \subset \mathbb{R}^2$  be a bounded domain with smooth boundary  $\partial \Omega$ .

- 1. If  $\Omega$  is the unit disk,  $\alpha \chi ||u_0||_{L^1(\Omega)} < 8\pi k_c$  and  $u_0(x) = u_0(-x)$ ,  $v_0(x) = v_0(-x)$  hold, then the solution of (10) exists globally in time and satisfies the equations in the classical sense, i.e. the solution is sufficiently smooth.
- 2. If  $T_{\max} < \infty$  then

 $\lim_{t \to T_{\max}} ||u(t) \log u(t)||_{L^{1}(\Omega)} = \lim_{t \to T_{\max}} ||u(t)v(t)||_{L^{1}(\Omega)} = \infty$ 

and

$$\lim_{t \to T_{\max}} ||\nabla v(t)||^2_{L^2(\Omega)} = \lim_{t \to T_{\max}} \int_{\Omega} e^{av(t)} dx = \infty,$$
  
where  $a > 1$ .

3. If  $\Omega$  is simply connected,  $\alpha \chi ||u_0||_{L^1(\Omega)} < 8\pi k_c$ , and  $T_{\max} < \infty$  then

$$\lim_{t\to T_{\max}}\int_{\partial\Omega}e^{\nu(t)/2}dx=\infty.$$

The last statement in particular implies together with the previous statements on the number of isolated blow-up points in Theorem 10 that if there is a solution that blows up in finite time for  $4\pi k_c < \alpha \chi ||u_0||_{L^1(\Omega)} < 8\pi k_c$  then the blow-up has to happen at the boundary of the domain. However, at this stage it has to be pointed out that these results do not give the existence of a solution that blows up in either finite or infinite time. These results always use the existence of a solution that blows up in finite time as an assumption, but do not prove that those solutions in fact really exist.

Beside the analysis of the Keller-Segel system (10) on a bounded domain T. Nagai also studied the problem on the whole space  $\mathbb{R}^2$ . In this situation he could prove that for  $\alpha \chi \int u_0(x) dx < 4\pi k_c$  the solution exists globally in time, once again via analyzing  $\mathbb{R}^2$ 

the functional F(u, v) for  $\Omega = \mathbb{R}^2$  this time. Furthermore he found several decay properties of the solution but for those results I refer the reader to [99].

Throughout this subsection we focused on the two dimensional case and left out the other space dimensions. So what is known for the cases N = 1 and  $N \ge 3$ ? For the case N = 1 the paper by K. Osaki and A. Yagi [113] fills the gap of the missing global existence proof for (10). Furthermore they show there that the  $\omega$ -limit set of the solution contains at least one stationary solution. For the case of higher space dimensions  $N \ge 3$  and a bounded domain  $\Omega \subset \mathbf{IR}^N$  I am aware of any result on the time asymptotic behaviour of the solution. The local existence of a solution can be established in such cases
using the results of H. Amann [6, 7] for example. This has been mentioned for example in [62] and [119].

### 4.3.2 Results for system (15)

Independent from the previous line of research and parallel to those results there were the results for system (15). Under different regularity assumptions on the domain and the solution than those assumed in [94] and [158] H. Gajewski and K. Zacharias proved in [44] the local existence of a weak solution of (15) where they defined a weak solution of (15) in the following way.

**Definition 2 (Gajewski & Zacharias)** A pair of functions (U(t, x), V(t, x)) with

$$\begin{split} & U \in L^{\infty}(0,T;L^{\infty}_{+}(\Omega)) \cap L^{2}(0,T;H^{1}(\Omega)), \quad U_{t} \in L^{2}(0,T;(H^{1}(\Omega))^{*}), \\ & V \in L^{\infty}(0,T;L^{\infty}(\Omega)) \cap C(0,T;H^{1}(\Omega)), \qquad V_{t} \in L^{2}(0,T;L^{2}(\Omega)) \end{split}$$

is called a weak solution of (15) if for all  $h \in L^2(0, T; H^1(\Omega))$  the following identities hold:

$$0 = \int_{0}^{T} \langle U_{t}, h \rangle \, dt + \int_{0}^{T} \int_{\Omega} (\nabla U - U \nabla V) \cdot \nabla h \, dx \, dt,$$
  
$$0 = \int_{0}^{T} \int_{\Omega} V_{t}h \, dx \, dt + \int_{0}^{T} \int_{\Omega} (k_{c} \nabla V \cdot \nabla h + (\gamma V - \alpha \chi (U - 1)) \cdot h) \, dx \, dt.$$

Their existence result is:

**Theorem 12 (Gajewski & Zacharias)** Let  $\Omega \subset \mathbb{R}^2$  be a bounded domain and its boundary is piecewise from the class  $C^2$ . For  $U_0 \in L^{\infty}_+(\Omega)$  and  $V_0 \in W^{1,p}(\Omega)$ , p > 2, and appropriate T > 0 there is a unique weak solution of (15) with  $U(0) = U_0$ ,  $V(0) = V_0$ . Moreover, for  $0 \le t < T$  it holds  $t \mapsto U(t) \in L^{\infty}_+$  and the function  $t \mapsto ||\nabla V(t)||^2_{L^2(\Omega)}$  is absolutely continuous on [0, T].

For (15) the Lyapunov function F takes the following form:

$$F(U(t), V(t)) := \frac{1}{2\alpha\chi} \int_{\Omega} k_c |\nabla V(t)|^2 + \gamma V^2(t) \, dx + \int_{\Omega} U(t)(\log(U(t)) - 1) + 1 - (U(t) - 1)V(t) dx.$$

In fact Gajewski and Zacharias showed that one can bound F by a functional only depending on V, namely

$$F(U(t), V(t)) \ge \mathcal{F}(V(t)) = \frac{1}{2\alpha\chi} \int_{\Omega} k_c |\nabla V(t)|^2 + \gamma V(t)^2 dx - |\Omega| \log\left(\frac{1}{|\Omega|} \int_{\Omega} e^{V(t)} dx\right).$$

Using the Moser-Trudinger type inequality by Chang and Yang in [23] it is possible to show that  $\mathcal{F}(V)$  is lower semicontinuous and coercive on the set  $\mathcal{D} := \{V \in H^1(\Omega) \mid V \text{ has mean value zero over the domain } \Omega\}$  if  $\alpha \chi |\Omega| < 4\Theta k_c$ , where  $\Theta$  denotes the smallest interior angle of the piecewise smooth domain  $\Omega$ . Therefore the calculus of variations guarantees the existence of a minimizer of  $\mathcal{F}$  over the set  $\mathcal{D}$ . As a conclusion we get the

	775N 2015 731			
Ibr	roio	htor	ortil	0
111	-ISIC:	11150	1111K	E

boundedness of the functional F from below in this case. The boundedness of the Lyapunov functional and the fact that for

$$U(x) = \frac{|\Omega|e^{V(x)}}{\left(\int_{\Omega} e^{V(x)} dx\right)}$$

the equality  $F(U, V) = \mathcal{F}(V)$  holds lead Gajewski and Zacharias to:

**Theorem 13 (Gajewski & Zacharias)** Let  $\alpha \chi |\Omega| < 4\Theta k_c$ . Then there exist a sequence  $t_k \to \infty$  and functions  $U^*$ ,  $V^*$  such that

$$U(t_k) \rightarrow U^*$$
 in  $L^2(\Omega), V(t_k) \rightarrow V^*$  in  $H^1(\Omega),$ 

and

$$F(U(t_k), V(t_k)) \to F(U^*, V^*)$$
 as  $t_k \to \infty$ .

Moreover the identity

$$U^* = \frac{|\Omega| e^{V^*}}{\left(\int\limits_{\Omega} e^{V^*} dx\right)}$$

holds, and  $V^*$  is the solution of the boundary value problem

$$-k_c \Delta V^* + \gamma V^* = \alpha \chi (U^* - 1) \text{ in } \Omega, \ \frac{\partial V^*}{\partial n} = 0 \text{ on } \partial \Omega.$$

As it has been shown in [63, Theorem 3, page 408] the previous result does not only hold for subsequences. Furthermore the steady state might also be nontrivial in the case where  $\alpha \chi |\Omega| < 4\Theta k_c$ . Gajewski and Zacharias presented in [44, Proposition 5.3, page 109] an example in which  $\Omega := \{(x, y) : 0 < x < a, 0 < y < b\}$  denotes a rectangle where

$$ab < \frac{2\pi k_c}{\alpha \chi}$$
 and  $a^2 > \frac{\pi^2 k_c}{\alpha \chi (\log(4) - 1) - \gamma} > 0$ 

and the initial data  $(U_0(x), V_0(x))$  is given by

$$U_0(x) = 1 + \cos\left(\frac{\pi x}{a}\right)$$
 and  $V_0(x) = \cos\left(\frac{\pi x}{a}\right)$ .

We then see that

 $F(U_0, V_0) < 0 = F(1, 0)$ 

and thus there has to be a nontrivial stationary solution of system (15) also in this case and not only in the cases mentioned in Section 3.

The boundedness of the Lyapunov functional F(U, V) by the functional  $\mathcal{F}(V)$  has several consequences that are demonstrated in [63]. However using the same sequence as in the proof of the existence of a nontrivial steady state solution in section 3 one can show for  $\alpha \chi |\Omega| > 4k_c \pi$  there is a sequence of functions  $\{(U_{\varepsilon}, V_{\varepsilon})\}_{\varepsilon>0}$  such that

$$F(U_{\varepsilon}, V_{\varepsilon}) \to -\infty$$
 as  $\varepsilon \to 0$ .

Furthermore, if  $\Omega \subset \mathbb{R}^2$  is simply connected and if  $\alpha \chi |\Omega| < 8\pi k_c$ ,  $p \in (1, 8\pi k_c/\alpha \chi |\Omega|)$  is arbitrary but fixed and q > 1 is such that 1 = 1/p + 1/q then one can bound the functional  $\mathcal{F}(V)$  from below in the following way:

$$\mathcal{F}(V) \geq \int_{\Omega} \left( \frac{k_c}{2\alpha\chi} - \frac{p|\Omega|}{16\pi} \right) |\nabla V|^2 + \frac{\gamma}{2\alpha\chi} V^2 \, dx - \frac{2|\Omega|}{q} \log \left( \int_{\partial\Omega} e^{qV/2} \, dS \right) + K(p, q, \alpha, \chi, k_c, |\Omega|),$$

where  $K(p, q, \alpha, \chi, k_c, |\Omega|)$  denotes a constant depending on  $p, q, \alpha, \chi, k_c$  and  $|\Omega|$ . However this has the consequence that we have the following blow-up result which summarizes the results from [61, 63, 64, 65].

**Theorem 14 (Horstmann & Wang)** Let  $\Omega \subset \mathbb{R}^2$  be a smooth, simply connected domain and  $\gamma > 0$ . Furthermore assume that

 $4k_c\pi < \alpha\chi|\Omega|$  and that  $\alpha\chi|\Omega|/k_c \neq 4\pi m$  for  $m \in \mathbb{N}$ ,

then there exist a constant  $-\infty < \hat{K} \le 0$  and initial data  $(U_0, V_0)$ , such that

 $\hat{K} > F(U_0, V_0)$ 

and the corresponding solution of (15) blows up in finite or infinite time. For these blow-up solutions the following statements hold:

$$I. \lim_{t \to T_{\max}} ||U(t, \cdot)||_{L^{2}(\Omega)} = \lim_{t \to T_{\max}} ||U(t, \cdot)||_{L^{\infty}(\Omega)} = \infty$$
$$2. \lim_{t \to T_{\max}} ||U(t) \log U(t)||_{L^{1}(\Omega)} = \lim_{t \to T_{\max}} \int_{\Omega} U(t, x) V(t, x) \, dx = \infty$$

$$3. \lim_{t \to T_{\max}} ||\nabla V(t, \cdot)||_{L^2(\Omega)} = \lim_{t \to T_{\max}} \int_{\Omega} e^{V(t, x)} dx = \lim_{t \to T_{\max}} ||V(t, \cdot)||_{L^{\infty}(\Omega)} = \infty$$

4. If  $4\pi k_c < \alpha \chi |\Omega| < 8\pi k_c$  and  $\Omega$  is a simply connected domain, then

$$\lim_{t \to T_{\max}} \int_{\partial \Omega} e^{qV(t,x)/2} dS = \infty$$

for every 
$$q \in (8\pi k_c/(8\pi k_c - \alpha \chi |\Omega|), \infty)$$
.

There are technical reasons why one has to exclude the multiples of  $4\pi$  in the theorem. From the biological point of view this makes no sense and there is a hint in [70] that in Theorem 14 the statements are in fact true if one only assumes  $4k_c\pi < \alpha\chi|\Omega|$ . For  $\gamma = 0$  there is a similar result for radially symmetric initial data in [65]. Since this proof is easy to illustrate I give this theorem and the sketch of the proof, too. So we can formulate our blow-up result.

**Theorem 15 (Horstmann)** Let  $\Omega = B(0, R) \subset \mathbb{R}^2$ . Further assume that  $\gamma = 0$ ,  $8k_c\pi < \alpha\chi|\Omega|$  and  $\alpha\chi|\Omega|/k_c \neq 8\pi q$ ,  $q \in \mathbb{N}$ ,

l'Ibersichtsartikel	Historischer Artikel	Buchhesprechungen
Obersichtsaltikei	historischer Artiker	Duchbesprechungen

then there exist radially symmetric initial data  $(U_0, V_0)$  and a constant  $\hat{K}$ , such that

$$K > F(U_0, V_0)$$

and the corresponding solution of (15) blows up in finite or infinite time.

The proof of Theorem 15 is easily demonstrated. First one shows the existence of the constant  $\hat{K}$  via contradiction. Thus one assumes that there is no such constant. Therefore there exists for  $\Omega = B(0, R) \subset \mathbb{R}^2$  a sequence  $(v_m)_{m \in \mathbb{N}} \in \mathcal{D}$  of solutions of the equation

$$\begin{cases} -k_c \Delta v_m = \alpha \chi \left( |\Omega| e^{v_m} \middle/ \int_{\Omega} e^{v_m} dx - 1 \right), & \text{in } \Omega \\ \partial v_m / \partial n = 0, & \text{on } \partial \Omega, \end{cases}$$
  
with  $\int_{\Omega} |\nabla v_m|^2 dx < \infty \ \forall m \in \mathbb{N}, \ \lim_{m \to \infty} \int_{\Omega} |\nabla v_m|^2 dx = \infty$ 

and

$$\lim_{m\to\infty}\log\left(\frac{1}{|\Omega|}\int\limits_{\Omega}e^{\nu m}dx\right)=\infty.$$

These sequence can therefore be identified as a sequence of stationary solutions of system (15) in the radially symmetric setting and with  $\gamma = 0$ . Using the transformation

$$w_m = v_m - \log\left(\frac{1}{|\Omega|}\int\limits_{\Omega} e^{v_m}\right) - \frac{\alpha\chi}{4k_c}|x|^2$$

the function  $w_m$  solves the problem

$$\begin{cases} -\Delta w_m = \frac{\alpha\chi}{k_c} e^{w_m + (\alpha\chi/4k_c)|x|^2}, & \text{in } \Omega\\ \partial w_m/\partial n = -\frac{\alpha\chi}{2k_c} (x \cdot n(x)), & \text{on } \partial\Omega, \end{cases}$$
  
with  $\int_{\Omega} |\nabla w_m|^2 dx < \infty \ \forall m \in \mathbb{N}, \ \lim_{m \to \infty} \int_{\Omega} |\nabla w_m|^2 dx = \infty$ 

and

$$\frac{1}{|\Omega|} \int_{\Omega} e^{w_m + \frac{\alpha \chi}{4k_c} |x|^2} dx = 1 \ \forall m \in \mathbb{N}.$$

Using the results from [78] and the Sobolev imbedding theorems we see that  $v_m$  is in fact  $C^{2,\beta}(\overline{\Omega})$  provided  $\partial\Omega$  is Lipschitz and thus  $w_m$  also belongs to  $C^{2,\beta}(\overline{\Omega})$ . According to a result from Brézis and Merle [20] there exists a subsequence  $(w_{m_i})_{m_i \in \mathbb{N}}$  for these  $(w_m)_{m \in \mathbb{N}}$  such that one of the following three alternatives holds:

- 1. The sequence  $(w_{m_i})_{m_i \in \mathbb{N}}$  is uniformly bounded in  $L^{\infty}_{loc}(\Omega)$ .
- 2. For each compact subset  $\mathcal{K} \subset \Omega$  we have:  $\sup_{k} w_{m_i} \to -\infty$  uniformly, as  $m_i \to \infty$ .
- 3. There exists a blow-up set  $\mathcal{BS} = \{p_1, ..., p_m\} \subset \Omega$  and sequences  $(x_{m_i}^j)_{j \in \{1,...,m\}} \subset \Omega$  such that for  $m_i \to \infty$ ,  $x_{m_i}^j \to p, w \in (x_i^j) \to \infty$  for i = 1, m.

$$x_{m_i}^j \to p_j, \ w_{m_i}(x_{m_i}^j) \to \infty \text{ for } j = 1...m.$$

Furthermore, on each compact subset  $\mathcal{K} \subset \Omega \setminus \mathcal{BS}$  we have

$$\sup_{\mathcal{K}} w_{m_i} \to -\infty, \text{ as } m_i \to \infty$$

and

$$\frac{\alpha \chi}{k_c} e^{w_{m_i} + (\alpha \chi/4k_c)|x|^2} \to \sum_{j=1}^m 8\pi q_j \delta_{x=p_j}$$

in the sense of measure, where  $q_j \in \mathbb{N}$ . (See [82] for the statement about the  $q_j$ .)

However as it has been done in [65] one can show that none of these alternatives is possible for such a sequence of stationary solutions. Therefore such a sequence cannot exist and one can conclude that there exists a constant  $\hat{K} \in \mathbb{R}$  ( $\hat{K} \leq 0$ ), such that for all radially symmetric stationary solutions (U, V) of system (15)

 $F(U,V) \ge \hat{K} > -\infty$ 

holds. Now let us choose a  $\varepsilon_0$  arbitrary but fixed, such that  $\hat{K} > \mathcal{F}(V_{\varepsilon_0}(x))$  where

$$V_{\varepsilon_0}(x) = \log\left(\frac{\varepsilon_0^2}{(\varepsilon_0^2 + \pi |x|^2)^2}\right) - \frac{1}{|\Omega|} \int_{\Omega} \log\left(\frac{\varepsilon_0^2}{(\varepsilon_0^2 + \pi |x|^2)^2}\right) dx.$$

We see that  $V_{\varepsilon_0}(x) \in W^{1,\infty}(\Omega)$ . Now let us set

$$U_{arepsilon_0}(x) = rac{|\Omega| e^{V_{arepsilon_0}(x)}}{\int\limits_{\Omega} e^{V_{arepsilon_0}}(x) dx}.$$

We see that  $U_{\varepsilon_0} \in L^{\infty}_+(\Omega)$  and that

$$F(U_{\varepsilon_0}(x), V_{\varepsilon_0}(x)) = \mathcal{F}(V_{\varepsilon_0}(x)) < \hat{K}.$$

Choosing  $U_0(x) = U_{\varepsilon_0}(x)$  and  $V_0(x) = V_{\varepsilon_0}(x)$  the corresponding solution of the Keller-Segel model (15) has to blow up in finite or infinite time.

Of course there are questions directly connected with the above results. The two most important are:

1. Is the blow-up time for the blow-up solution from Theorem 14 and Theorem 15 finite?

Obersichtsattiker – Historischer Aftiker – Duchbesprechu	Übersichtsartikel	Historischer Artikel	Buchbesprechung
--	-------------------	----------------------	-----------------

2. Suppose  $T_{\max} < \infty$ , does either  $\inf_{0 \le t < T_{\max}} F(U(t), V(t)) > -\infty$ or  $\lim_{t \to T_{\max}} F(U(t), V(t)) = -\infty$  hold?

H. Gajewski and K. Zacharias gave in [44, p. 94 & 95] an example for initial data for system (15) such that the solution blows up in the corner of a rhombic domain. They considered the domain

$$\Omega = \left\{ (x, y) \mid \frac{|x|}{a} + \frac{|y|}{b} < 1, \ a = \sqrt{\frac{\tan(\Theta/2)}{2}}, \ b = \frac{1}{\sqrt{2\tan(\Theta/2)}} \right\}$$

with an acute opening angle  $\Theta \leq \pi/2$ . For  $\alpha = \chi = k_c = 1$  they used the initial data

$$u_0(x) = U_0(x)/\overline{U_0(x)},$$

where

$$U_0(x) = \frac{8(1+\sigma)}{\sigma} \exp\left(-\frac{|x|}{\sigma}\right)$$

with  $0 < \sigma < 1$  and as  $V_0(x)$  the solution of the boundary value problem

$$\Delta V_0 + \overline{U_0}(u_0 - 1) = 0$$
 in  $\Omega$ ,  $\frac{\partial V_0}{\partial n} = 0$  on  $\partial \Omega$ .

The corresponding solution (U(t), V(t)) of the equations (15) blows up in finite time in a corner of the domain. Furthermore their numerical calculations showed that for this solution the Lyapunov functional  $F(U(t), V(t)) \rightarrow -\infty$  in finite time. There is also another numerical example given in [43] where the initial data is such that the function u(x,0) already has its maximum in the corner with the smallest interior angle of the rhombic domain. The solution then blows up in this corner in finite time.

Some known results with their references are summarized once again in table 5.

Dimension	Observation	References
N = 1	The solution of the Keller-Segel model exists globally in time and for any initial data $u_0 \in L^2(\Omega)$ , $v_0(x) \in H^1(\Omega)$ with $u_0 \ge 0$ , $\int_{\Omega} u_0(x) dx = M < \infty$ , $\inf_{x \in \Omega} v_0(x) > 0$ the $\omega$ -limit set of the solution contains at least one stationary solution.	[113]
<i>N</i> = 2	If $\alpha \chi \int_{\Omega} u_0(x) dx < 4\pi k_c$ , then the solution exists globally in time and its $L^{\infty}$ -norm is uniformly bounded for all times. Furthermore it converges to a stationary solution as $t \to \infty$ .	[11, 44, 63] and [94]
	If $4\pi k_c < \alpha \chi \int_{\Omega} u_0(x) dx < 8\pi k_c$ , then there exist initial data such that the corresponding solution of the Keller-Segel model blows up at the boundary of $\Omega$ either in finite or in infinite time.	[63, 64, 135]
	If $8\pi k_c < \alpha \chi \int_{\Omega} u_0(x) dx$ , then there exist initial data such that the corresponding solution of the Keller-Segel model blows up either in finite or in infinite time.	[52, 61, 64] and [65]
	Furthermore there exist radially symmetric initial data such that $u(t, x)$ forms a $\delta$ -singularity in finite time in the center of a disk $\Omega$ .	[52]
	Given a blow-up solution and an isolated blow-up point $x_0$ , we have $0 < R << 1, m \ge m^*$ , and $f \in L^1(B(x_0, P) \cap \Omega) \cap C(\overline{B(x_0, R) \cap \Omega} \setminus \{x_0\})$ satisfying $f \ge 0$ and $u(t, \cdot)dx$ converges weakly to $m\delta_{x_0}(dx) + f dx$ as $t \to T_{finite}$ in the set of Radon measures on $\overline{B(x_0, R) \cap \Omega}$ , where $m^*$ is either $8\pi$ , for $x_0 \in \Omega$ or $4\pi$ for $x_0 \in \partial\Omega$ .	[100]
	If the blow-up time is finite and (26) holds there exist only isolated blow-up points.	
	If (10) is radially symmetric and $T_{\max} < \infty$ then the set of blow-up point consists only of the origin $\{0\}$ .	
	If $\Omega = \mathbb{R}^2$ and $\alpha \chi \int_{\mathbb{R}^2} u_0(x) dx < 4\pi k_c$ then the solution of (10) exists globally in time.	[99]
<i>N</i> = 3	Let $\Omega$ be a smoothly bounded domain in $\mathbb{R}^3$ . For sufficiently smooth initial data, satisfying the boundary data there exists a unique solution of (10) locally in time. Furthermore for all $T > 0$ there exists a constant $C_T$ , such that if the initial data satisfies $  v_0  _{H^2(\Omega)} < C_T$ , $  u_0  _{L^{\infty}(\Omega)} < C_T$ and $  \nabla u_0  _{L^2(\Omega)} < C_T$ , then the problem (10) has a unique solution on $[0, T] \times \Omega$ .	[16]

Table 5: Possible time asymptotical behaviour of the solutions of (10) and of (15).

#### 4.4 Results for related systems

There are some related results that should be mentioned at this point of this survey. P. Biler studied in [11] system (10) with different boundary conditions. For system (10) his local existence result is pretty much the same as the local existence result of Gajewski and Zacharias [44].

However for the system

(27) 
$$\begin{cases} u_t = \nabla(\nabla u - \chi u \nabla v), & x \in \Omega, \ t > 0\\ 0 = k_c \Delta v - \gamma v + \alpha u, & x \in \Omega, \ t > 0\\ 0 = \partial u / \partial n - \chi u \partial v / \partial n, \ x \in \partial \Omega, \ t > 0, \quad v(t, x) = K_{\gamma} * (\alpha u(t, x)), \\ u(0, x) = u_0(x), & x \in \Omega, \end{cases}$$

where  $K_{\gamma}$  denotes the Bessel potential, Biler proves the following finite time blow up result:

**Theorem 16 (Biler)** If  $\Omega \subset \mathbb{R}^N$ ,  $N \ge 2$ , is a bounded star-shaped domain (with respect to the origin), then for  $u_0(x)$  with sufficiently large  $||u_0||_{L^1(\Omega)} = M$ , there is no global in time solution of (27).

Results similar to those results in [64, 65] have been proven by G. Wolansky in [155] for the system

(28) 
$$\begin{cases} 0 = \nabla(\nabla u - \chi u \nabla v), & x \in \Omega, \ t > 0 \\ v_t = k_c \Delta v + \alpha u, & x \in \Omega, \ t > 0 \\ 0 = \partial u / \partial n - \chi u \partial v / \partial n, \ v(t, x) = 0, & x \in \partial \Omega, \ t > 0 \\ u(0, x) = u_0(x), \ v(0, x) = v_0(x) & x \in \Omega. \end{cases}$$

This system contains an elliptic equation for the myxamoebae density and a parabolic equation for the cAMP-concentration similar to the second equation in (10). Even though he also has a Lyapunov function the techniques he used to prove his blow up result can only be applied to a system with Dirichlet boundary conditions for the second equation. They fail in the case of Neumann boundary conditions as they are treated in [64, 65]. In [155] G. Wolansky is led to an equation for the stationary solutions of his model which is similar to problem (13). However in his case the equation is equipped with Dirichlet boundary data, which allows him to use different arguments (more precisely the moving plane method see [31, pp. 521–522]) to exclude the first alternative of the Brézis and Merle [20] result. In the case that is mentioned in the previous section and [64, 65] one has to use different techniques to get rid of the possible alternatives stated by Brézis and Merle in [20].

# 5 Comparison of the questions asked by Jäger and Luckhaus with the results so far

Let us now take some time and let us see which questions of those asked in [69] have been answered up to now and which remain open. W. Jäger and S. Luckhaus asked about more information on the set of blow-up points. We have seen in the previous sections that there exists the possibility of blow-up points in the interior and at the boundary of a domain  $\Omega \subset \mathbb{R}^2$ . Also the upper bound of the possible number of blow-up points is sharp and known, however a better lower bound is still needed. Also the location of the boundary blow-up points should be studied more carefully. For smooth domains the boundary blow-up point should be a point of maximal mean curvature which would correspond with the numerical calculations of H. Gajewski and K. Zacharias in [44] for piecewise smooth domains and with the results and hints from the steady state analysis resp. the shape of the least energy solutions.

The question whether the solution exists globally in time as a weak solution can be negated. However it might be possible to study the problem for a different formulation of a solution like  $L^1$ -solutions. But as far as I know there has not been any attempt to do so up to now. For the third question we turn to an own subsection.

#### 5.1 What happens after blow-up?

In connection with the question "What happens after the blow up of the solution?" that was already asked in [69], we have seen that the solution does not exist globally in time as a weak  $H^1$ -solution, but is there a notation of a measure valued solution or  $L^1$ -solution for the Keller-Segel model? With such a notation, which would be natural since the solution belongs to  $L^1(\Omega)$  for all times, it would make sense to study the possible movement of the aggregation centers in the considered domain.

Using a different approach than the idea of introducing a new notation of a solution as just mentioned, J. J. L. Velázquez made the first step to give an answer to the question what will happen after blow-up in [151, 152].

In [150] J. J. L. Velázquez studied the question whether aggregation at the interior of the boundary of the domain  $\Omega \subset \mathbb{R}^2$  takes place in a stable manner, or, if on the contrary, solutions exhibit a tendency to move towards the boundary. His result is that after small pertubations of the solution found in [52], the new solution will blow up in a manner entirely similar but in a slightly shifted point of  $\Omega$  at a slightly different time. Thus his computations indicate that the possibility of aggregates with high density of *u* moving quickly towards the boundary does not exist.

He then studies in [151, 152] the system

(29) 
$$\begin{cases} u_t = \nabla(\nabla u - G_{\epsilon}(u)\nabla v), & x \in \mathbb{R}^2, \quad t > 0\\ 0 = \Delta v + u, & x \in \mathbb{R}^2, \quad t > 0, \end{cases}$$

where  $G_{\epsilon}(u) = \frac{1}{\epsilon} Q(\epsilon u)$  for a small parameter  $\epsilon > 0$  and an increasing function Q(s) satisfying  $Q(s) = s + O(s^2)$  as  $s \to 0$  and  $Q(s) \sim L$  as  $s \to \infty$ , where L > 0 is a given number. For  $Q(s) = \frac{s}{1+s}$  and  $\epsilon = 0$  the system becomes formally system (16). For  $\epsilon > 0$  the solution of (29) exist globally in time under general assumptions on the initial data. Thus it is a natural question to try to understand the asymptotics of the solution of (29) when  $\epsilon$  approaches zero. This has been done in [151] using asymptotic expansions. Using these methods Velázquez showed that solutions of system (29) exist, that have a finite amount of mass  $M_i(t)$  concentrated in a neighbourhood of a set of points  $x_i(t)$ .

Übersichtsartikel	Historischer Artikel	Buchbesprechungen
		Edonie ooproon angen

Velázquez studies the motion of those regions with high densities of u in [151] and derives a system of equations that describes the dynamics of these regions. According to the formal limit as  $\epsilon \to 0$  his results can somehow be interpreted as a formal explanation of the behaviour of the solution of (16) after blow up. The well-posedness of the derived system for the dynamics of the high density regions is established in [152].

## 6 More general formulations of the chemotaxis equations

The original formulation of the Keller-Segel model allowed more general functional forms than we assumed in the last section. Even though the question whether the given functional form represents the situation in Dictyostelium aggregation in an appropriate way should be discussed, the system is adequate to describe chemotactical movement of mobile species. A number of possible plausible functional forms has been proposed by E.F. Keller in [76]. Several of these proposed functional forms will also be discussed in the upcoming subsections. Furthermore she discussed the possible existence of traveling wave solutions, a topic which will be in the center of our observations later in this paper. Since there is a large number of different examples for species that move positive chemotactically and also a large variety of different models for the chemotactical sensing of the particular species (see for example [84] for a model of the cAMP production and sensing mechanism in Dictyostelium discoideum) it is useful to try to find a more general theory that contains a larger class of possible models. Let us see what results are available in this cases. So let us now turn to more general formulations of the system without having a particular biological example in mind. So we focus on the following system of two nonlinear parabolic partial differential equations, which is given by

(30) 
$$\begin{cases} u_t = \nabla(k(u,v)\nabla u - h(u,v)\nabla v), & x \in \Omega, \quad t > 0\\ v_t = k_c \Delta v - f(v)v + g(u,v), & x \in \Omega, \quad t > 0 \end{cases}$$

for  $\Omega \subset \mathbf{I}\!\mathbf{R}^N$  completed with either

- (31)  $\frac{\partial u}{\partial n} = \frac{\partial v}{\partial n} = 0 \text{ on } \partial \Omega \times \{t > 0\},$
- (32) or u = 0, v = 0 on  $\partial \Omega \times \{t > 0\}$ ,

(33) or 
$$k(u, v) \frac{\partial u}{\partial n} - h(u, v) \frac{\partial v}{\partial n} = 0, v = 0 \text{ on } \partial\Omega \times \{t > 0\}$$

as boundary conditions and initial data  $u(0, x) = u_0(x)$  and  $v(0, x) = v_0(x)$  for  $x \in \Omega$ . Here  $k_c$  is once again a positive constant. For the functions appearing in the model the following conditions have been considered to be reasonable: k(r, s) > 0 for all  $(r, s) \in \mathbb{R} \times \mathbb{R}$ , the function f satisfies  $f(s) \ge \text{const}$  for all  $s \in \mathbb{R}$  and  $\frac{\partial}{\partial r}g(r, s) \ne 0$  holds for all  $(r, s) \in \mathbb{R} \times \mathbb{R}$ .

The question whether a solution to such problems exist locally in time has been studied in [62] using results by H. Amann [6, 7] and in [158] using other techniques. As it was mentioned in the previous section a Lyapunov function is a helpful tool for analyzing the time asymptotic behaviour of the solution. Thus one wonders under which conditions the present system has a Lyapunov function. Therefore we turn to this question next.

#### 6.1 Lyapunov functions

For the rest of the present section we will use the following notations:

$$F(v) := \int_{-\infty}^{v} f(s)s \ ds \ and \ G(u,v) := -\int_{-\infty}^{v} g(u,s) \ ds.$$

At some places of the present section we will assume that

(34) 
$$\int_{\Omega} F(v) \, dx \ge k_1 \int_{\Omega} v^2 \, dx$$

is true, where  $k_1$  is a nonnegative constant (If we have homogeneous Neumann boundary data we assume  $k_1 > 0$ !).

**Theorem 17 (Horstmann)** If there exists a function R(u) such that

$$\frac{h(u,v)}{k(u,v)} \left[ \frac{\partial^2}{\partial u^2} G(u,v) + \frac{d^2}{du^2} R(u) \right] + \frac{\partial^2}{\partial u \partial v} G(u,v) = 0,$$

then there exists a Lyapunov function for system (30), provided  $\frac{\partial^2}{\partial u^2} G(u, v) + \frac{d^2}{du^2} R(u) \ge 0$ holds true for the solution of (30). In the case of boundary condition (32) we have to assume additionally that  $\frac{\partial}{\partial u} G(0,0) = 0 = \frac{dR}{du}(0)$ . The Lyapunov function for system (30) is then given by

$$\mathcal{H}(u(t), v(t)) := \int_{\Omega} \frac{k_c}{2} |\nabla v(t)|^2 + F(v(t)) + R(u(t)) + G(u(t), v(t)) dx.$$

A large number of examples is given in [62]. Let us here only give two examples for which a Lyapunov function  $\mathcal{H}(u, v)$  exists.

1. Let consider (30) with h(u, v) = u,  $g(u, v) = \frac{1}{2}u^2 e^{-v}$ , k(u, v) = 1, f(v) arbitrary. Then we have the Lyapunov function

$$\mathcal{H}(u(t), v(t)) := \int_{\Omega} \frac{k_c}{2} |\nabla v(t)|^2 + F(v(t)) + \frac{1}{2} u^2(t) e^{-v(t)} dx, \text{ i.e. } R(u) = u^2/2.$$

One can also find a whole class of other examples where a Lyapunov function exists. Let us suppose that we study system (30) together with (31). Let

$$h(u, v) = h_2(u)\phi(v) \text{ and } g(u, v) = \phi(v) \int h_1(s) \, ds,$$
  
$$k(u, v) = \tilde{k}(u) + \frac{h_2(u)}{h_1(u)} \left(\frac{d}{du}h_1(u)\right) \int^v \phi(s) ds$$

and f(v) be arbitrary. We see that there is a function R(u) such that

$$\frac{d^2}{du^2}R(u) = \frac{k(u,v)h_1(u)}{h_2(u)} + \left(\frac{d}{du}h_1(u)\right)\int \phi(s) \ ds.$$

Of course we see in this example that the right hand side has to be independent of v. If this is the case, then there exists a Lyapunov function  $\mathcal{H}(u, v)$  of the type given above, which is possibly unbounded from below. This example includes the systems studied in [44] and [119]. In [119] we have  $h(u, v) = u\phi(v)$  (with  $\phi(v) > 0$ ),  $g(u, v) = u\phi(v)$ , k(u, v) = 1 and f(v) = const > 0. Finally we get in this case  $R(u) = u \log(u)$ . For further results concerning some special cases of this type of systems see [119].

In fact this result allows to make statements for a larger class of nonlinearities in g(u, v) than those studied before for system (30) (as far as the author knows). Under certain additional assumptions one can now formulate results for the time asymptotic behaviour of the solution. Therefore we now make the following main assumption for the rest of this section.

#### Main assumption:

(35) 
$$\int_{\Omega} G(u,v) + R(u)dx \ge k_2 \int_{\Omega} |\nabla v|^2 dx + \text{const with } \frac{k_c}{2} + k_2 > 0.$$

In some special cases of (30) one can show that the solution of (30) converges to a possibly nontrivial steady state as  $t \to \infty$  (see [44, 62] and [119]). The results of W. Alt [3], R. Schaaf [128] and K. Post [119] concerning the Keller-Segel model in chemotaxis seem to indicate that such behaviour can also be expected in a more general setting. The following theorem summarizes our results on this aspect.

**Theorem 18 (Horstmann)** Suppose that (u(t), v(t)) is a weak solution of (30) and that (34) as well as our main assumption (35) is satisfied. Furthermore let either

1. 
$$\left|\frac{\partial^2}{\partial u^2} G(u,v) + \frac{d^2}{du^2} R(u)\right| / k(u,v) \le k_3 \text{ and } \frac{\partial}{\partial u} G(u,v) + \frac{d}{du} R(u) > k_4 \text{ or}$$
2. 
$$0 < \left[\frac{\partial^2}{\partial u^2} G(u,v) + \frac{d^2}{du^2} R(u)\right] \exp\left(\frac{\partial}{\partial u} G(u,v) + \frac{d}{du} R(u)\right) \le k_5 k(u,v) \text{ and}$$

$$\sqrt{\frac{k(u(t),v(t))}{\frac{\partial^2}{\partial u^2} G(u(t),v(t)) + \frac{d^2}{du^2} R(u(t))}} \in L^2(\Omega)$$
for all  $t \ge 0$ 

be true for the solution (u(t), v(t)) of (30). Let additionally f be Hölder continuous with Hölder exponent  $\beta \leq 1$  such that  $0 < \beta \leq 1$  if  $N \leq 3$  or  $\beta < 2/N$  if N > 3. Finally assume that  $|f(v)| \leq K_f$  for all  $v \in \mathbb{R}$ . Then there exist a sequence  $(t_k)_{k \in \mathbb{N}}$  and two functions  $v^*$  and  $g^*$  such that  $v(t_k) \rightarrow v^*$  in  $H^1(\Omega)$  resp. in  $H^1_0(\Omega)$ ,  $f(v(t_k))v(t_k) \rightarrow f(v^*)v^*$  in  $L^2(\Omega)$ and  $g(u(t_k), v(t_k)) \rightarrow g^*$  in  $L^2(\Omega)$ . Furthermore

$$\int_{\Omega} k_c \nabla v^* \nabla \varphi + f(v^*) v^* \varphi \ dx = \int_{\Omega} g^* \varphi \ dx$$

for all  $\varphi \in H^1(\Omega)$  (resp.  $\varphi \in H^1_0(\Omega)$ ). Finally we see that

$$\exp\left(\frac{-\left[\frac{\partial}{\partial u}G(u(t_k),v(t_k))+\frac{d}{du}R(u(t_k))\right]}{2}\right) \to \text{const}$$

in  $L^2(\Omega)$  if 1. holds and, respectively,

$$\exp\left(\frac{\left[\frac{\partial}{\partial u}G(u(t_k),v(t_k))+\frac{d}{du}R(u(t_k))\right]}{2}\right) \to \text{const}$$

in  $L^2(\Omega)$  if 2. holds.

The previous given first example satisfies assumption 1. of Theorem 18 while the systems studied in [44] and [119] satisfy assumption 2. of Theorem 18. The proof of this theorem goes along the line of the proof of Theorem 5.2 in [44, page 107] and can be found in detail in [62].

Furthermore one can also formulate certain conditions under which some  $L^{p}$ -estimates for the solution are possible. This has also been done in [62].

More general forms of the Keller-Segel model (4) have also been studied by A. Yagi in [158] for the case of two space dimensions and in the case of one spatial dimension by K. Osaki and A. Yagi in [113]. In [113] the authors study the Keller-Segel model (4) with  $k_1(u, v) = const$ ,  $uf(v) - k_3(v)v = \alpha u - \gamma v$  and  $k_2(u, v) = u\chi(v)$  where  $\chi(s)$  is a smooth function of  $s \in (0, \infty)$  satisfying

$$|\chi^{(i)}(s)| \leq const \cdot \left(s + \frac{1}{s}\right)^r$$
,

for  $0 < s < \infty$ , i = 0, 1, 2 with some positive constant and exponent *r*. In [113] they show that there exists a compact set of finite fractal dimension which attracts the solutions exponentially.

A. Yagi studied in [158] the two dimensional case of the Keller-Segel model under the assumptions that  $k_1(u, v) = c_0 + c_1u + c_2v$  with a positive constant  $c_0 > 0$  and nonnegative constants  $c_1$ ,  $c_2$ , and assuming that  $k_2(u, v) = u\chi(v)$  with  $0 \le \chi(s) \le b_0(1 + \frac{1}{s})$ , and that  $\chi(s)$ ,  $k_3(s)$ , f(s) are smooth functions of  $s \in \mathbb{R}^+$  satisfying  $0 \le f(s) \le b_1$ ,  $b_2 \le k_3(s) \le b_3(s^{p_0} + 1)$ , where  $b_0$ ,  $b_1$ ,  $b_2$ ,  $b_3$  are positive constants strictly larger than zero and the exponent  $p_0 > 0$ . For the initial data he assumed that  $u_0(x) > 0$  on  $\overline{\Omega}$ ,  $v_0(x) \ge \mu_0 > 0$  on  $\overline{\Omega}$  belong to  $H^{1+\epsilon_0}(\Omega)$  with some exponent  $0 < \epsilon_0 \le 1$  and a positive constant  $\mu_0$ . Using semigroup theory Yagi established the local in time existence of an unique, positive, classical solution in the same space as it has already been mentioned in the case of a linear sensitivity function in a previous section. Furthermore he determines blow-up norms of the maximal solution.

#### 6.1.1 Results on finite time blow-up

As it has been already mentioned in the section on the steady state solutions of the Keller-Segel model Nanjundiah's conjecture also contained a statement on the time asymptotical behaviour of the solution of the Keller-Segel model with a logarithmic chemotactical sensitivity function. Therefore we will first look at the results for this conjecture and related results. In this subsection we consider (4) with  $k_1(u, v) = 1$ ,  $uf(v) - k_3(v)v = \frac{1}{\varepsilon}(u-v)$ ,  $k_c = \frac{1}{\varepsilon}$  and  $k_2(u, v)$  either  $\chi^{\underline{u}}_v$  or  $\chi puv^{p-1}$  for p > 0. In the limiting case  $\varepsilon = 0$  it is easy to show in the same way as it has been done in the case of a linear chemotactical sensitivity function that the solution exists globally in time and that the  $L^{\infty}$ -norm of the solution is uniformly bounded for all times. Thus the interesting cases are once more the higher dimensional ones. So let us summarize these cases:

- 1. Let  $k_2(u, v) = u\phi(v)$  where  $\int \phi(s) ds$  is a smooth function with  $\phi(s) > 0$  for s > 0.
  - (a) Let N = 1, ε = 0 and ∫ φ(s)ds be smooth on (0,∞). Then the solution of the Keller-Segel model exists globally in time and is uniformly bounded. (See [98].)
- 2. Let  $k_2(u, v) = \chi p u v^{p-1}$  for p > 0.
  - (a) Let N = 2 and  $\varepsilon = 0$ . If  $0 , then the solution of the Keller-Segel model exists globally in time and is uniformly bounded. If <math>\Omega$  is a disk,  $u_0$  is radially symmetric,  $\int_{\Omega} u_0(x) |x|^2 dx$  is sufficiently small and p > 1, then the corresponding solution of (4) blows up in finite time. (See [93, 98, 130].)
  - (b) Let  $N \ge 3$  and  $\varepsilon = 0$ . If  $\Omega$  is a disk,  $u_0$  is radially symmetric,  $\int_{\Omega} u_0(x)|x|^{(N-2)p+2}dx$  is sufficiently small and p > 0, then  $T_{\max} < \infty$  and the corresponding solution of (4) blows up in finite time. (See [93, 98, 130].)
- 3. Let  $k_2(u, v) = \chi \frac{u}{v}$ .
  - (a) Let N = 2 and  $\varepsilon = 0$ . If  $\Omega$  is a disk,  $u_0$  is radially symmetric, then the solution is globally bounded in time. (See [93, 98, 130].)
  - (b) Let  $N \ge 3$  and  $\varepsilon = 0$ . If  $\Omega$  is a disk,  $u_0$  is radially symmetric and  $\chi < 2/(N-2)$ , then the solution is globally bounded in time. (See [93, 98, 130].)
  - (c) Let N = 2 and  $\varepsilon = 1$ . If  $\chi < 1$ , then the solution of (4) exists globally in time and for T > 0 there exists a constant  $C_T < \infty$  such that

$$\sup_{\substack{0 \le t \le T}} \left( ||u(t, \cdot)||_{L^{\infty}(\Omega)} + ||v(t, \cdot)||_{L^{\infty}(\Omega)} \right) < C_T.$$
  
(See [95].)

- (d) Let N = 2 and  $\varepsilon = 1$ . If  $\Omega$  is a disk, the initial data  $(u_0(x), v_0(x))$  is radially symmetric and  $\chi < 5/2$ , then the solution exists globally in time. (See [95].)
- (e) Let  $N \ge 3$  and  $\varepsilon = 0$ . If  $\Omega$  is a disk,  $u_0$  is radially symmetric,  $\int_{\Omega} u_0(x)|x|^2 dx$  is sufficiently small and  $\chi > 2N/(N-2)$ , then the solution of (4) blows up in finite time. (See [93, 98, 130].)

Results similar to (3c) have been obtained by K. Post studying (4) with  $k_1(u, v) = 1$ ,  $k_2(u, v) = u\phi'(v)$ ,  $uf(v) - k_3(v)v = u\phi'(v) - v$  and  $k_c = 1$ . For the precise details I refer the reader to [119].

Rascle and Ziti analyzed in [122] the system

$$\begin{cases} u_t = \nabla(\mu \nabla u - \chi u v^{-\beta} \nabla v), & x \in \Omega, \ t > 0\\ v_t = -kuv^m, & x \in \Omega, \end{cases}$$

where the constants  $\chi, k > 0$ . They constructed self-similar solutions for this system assuming that  $m < \beta = 1$ . For  $\mu = 0$  and one space dimension they observed that the bacterial density concentrated in finite time at the origin. For two space dimensions and initial data for the bacterial density which is zero at the origin they derived chemotactic rings concentrated around the origin after finite time. In higher space dimensions they achieved blow-up of the solution by an initial singularity of the chemoattractant in the origin.

For  $\mu > 0$  Rascle and Ziti observed in one space dimension that there are smooth initial data leading to finite time blow-up of the solution, while they were unable to construct self-similar solutions in space dimension larger or equal to two for reasonable initial conditions.

For m = 1 and  $v^{-\beta}$  replaced by a more general sensitivity function  $\chi(v)$  satisfying  $\chi(s) > 0$  for  $s \in \mathbb{R}_+$  and

$$\inf_{s\geq 0}\frac{s\chi'(s)}{\chi(s)} > -1,$$

this system has also been studied by Corrias et al in [27]. They considered the case of a bounded domain and the case of  $\Omega = \mathbf{IR}^N$ . In both cases they proved an existence result and studied the behaviour of the solution as  $t \to \infty$ . For a bounded domain they showed that

$$u(t,\cdot) \rightarrow \frac{1}{|\Omega|} \int_{\Omega} u_0 \ dx \ \text{in } L^1(\Omega) \ \text{and} \ v(t,\cdot) \rightarrow 0 \ \text{in } L^p(\Omega), \ p < \infty.$$

Furthermore they show the existence of radially symmetric self-similar solutions for N = 2.

## 6.1.2 Prevention of overcrowding

There are different points of view whether blow-up in chemotaxis is relevant or not. In fact for the chemotaxis system introduced and derived in the Davis' case by Othmer and Stevens for a single particle in [116] blow-up in finite time seems to correspond with the fact that the particle is trapped respectively localizes in finite time at one particular place (see also [139] for more comments on that aspect). Thus blow-up really makes sense for their model. Furthermore, blowing up of the solution only describes a high concentration of the particle populations in some aggregation centers. (Of course the convergence of the solution to non-trivial steady state solutions can also describe aggregation, but they are not the only possibility.) Since the Keller-Segel model only wants to describe the aggregation phase of chemotactical movement and not the formation of a fruiting

1 ***	1				
	hore	10	ntea	rti	101
U	DEIS	101	noa	1 U I	NCI

body the blow-up question is surely worth studying. As J. J. L. Velázquez wrote in [151, pp. 1–2]:

"Blow-up usually takes place in physical or biological models if they are approximations of more realistic models, usually containing small parameter (say  $\epsilon > 0$ ), that cannot exhibit singular behaviours unless this parameter is set to zero. Suppose that for  $\epsilon = 0$  the limit problem can develop singularities in finite time. The behaviour of the complete model for  $\epsilon > 0$  usually is similar to that of the limit model away from the singularities. However, the features of the problem with  $\epsilon > 0$ but small are usually very different from that of the limit problem near the singularities. The presence of blow-up just indicates that the approximations that lead to the simpler model where blowup takes place are not valid anymore near the singularity and that the whole dynamics of the complete model needs to be taken into account here."

However there are also other models of Keller-Segel type which exclude the possibility of blow-up solutions directly by introducing some mechanisms, that provide to strong aggregations or where the chemical production and decay directly is such that blow-up is impossible. For example in the case of the linear chemotactical sensitivity function  $g(u(t, x), v(t, x)) \in L^4(\Omega)$  for all  $t \ge 0$  with an uniform bound for all  $t \ge 0$  guarantees the global existence of the solution of system (30) (see for example [62]). One model containing a prevention of an overcrowding of the chemotactical species has been proposed by T. Hillen and K. J. Painter in [57]. They considered system (30) on a  $C^3$ -differentiable, compact Riemannian manifold  $\mathcal{M}$  under the assumptions that k(u, v) = 1.  $h(u, v) = u\beta(u)\chi(v)$  where  $\beta, \chi$  are three times continuous differentiable functions satisfying  $\chi > 0$ ,  $\beta(0) > 0$  and there exists a  $\overline{u} > 0$  such that  $\beta(\overline{u}) = 0$  and  $\beta(u) > 0$  for  $0 < u < \overline{u}$ . They assumed that the function  $f(v) \equiv 0$  and that g(u, v) = 0 $g_1(u, v)u - g_2(u, v)v$  is twice continuously differentiable with a bounded death rate  $g_2 \ge \delta > 0$  and a birthrate  $g_1 \ge 0$ . In their paper they prove the global existence of the solution in this case and present numerical simulations for the time evolution of the system in one and two space dimensions. They also show the potential pattern variety of the final steady state patterns for their version of model (30).

#### 6.1.3 Chemotaxis equations with population growth

An extremely large number of models describing chemotactical movement for species in a reproduction stage can be found in the literature. In general they are based on some version of the Keller-Segel equations with an additional growth term in the first equation. For example A. Bonami, D. Hilhorst, E. Logak and M. Mimura consider in [13, 14] the following versions of the classical model:

(36) 
$$\begin{cases} u_t = \nabla(k_1(u)\nabla u - u\nabla\chi(v)) + f(u), & x \in \Omega, \ t > 0\\ \varepsilon v_t = k_c \Delta v - \gamma v + \alpha u, & x \in \Omega, \ t > 0 \end{cases}$$

where  $k_1(u) = 1$  and f(u) = u(1-u)(u-a) with a constant 0 < a < 1 (see also [49] for results related to this system). However also different functional forms for f(u) are thinkable. For example in [17] one finds the proposed functional form f(u) = au for a positive constant a > 0 and in [29] system (36) is studied with  $k_1(u) = u^m$  and  $f(u) = u(1 - u^p)$  with m > 1 and  $p \ge 1$ . Some effects of such growth terms on the various possible patterns that one can observe during the evolution of the solution will be mentioned in Section 8.5 of the present paper. For some results on a more general chemotaxis growth model we refer the interested reader to [154]. There X. Wang studied in one space dimension the steady state solutions of the system

(37) 
$$\begin{cases} u_t = \nabla(\lambda \nabla u - \chi u \nabla \phi(v)) + (kf(u) - \theta - \beta v)v, & x \in [0, 1], t > 0\\ v_t = \Delta v - f(u)v, & x \in [0, 1], t > 0\\ u_x = \chi u(\phi(v))_x \text{ at } x = 0, 1, \\ v_x(0) = 0, v_x(1) = h(1 - u(1)), \end{cases}$$

where  $\lambda$ , k,  $\theta$ , h and  $\beta$  are positive constants and  $\chi \ge 0$  for different possible growth terms  $f(u) \in C^3([0,\infty))$  and chemotactic sensitivity functions  $\phi(v) \in C^5([0,\infty))$  satisfying f(0) = 0, f'(s) > 0 and  $\phi'(s) > 0$  for  $s \in [0,\infty)$ . He also proves the global existence and boundedness of the solution for those different growth factors for the population density.

## 7 The comparison principles by W. Alt for chemotaxis equations

In his (unfortunately almost unknown) Habilitation [3] from 1980 Wolfgang Alt studies quasilinear parabolic and elliptic systems including the chemotaxis equations by Keller and Segel with and without growth terms and for single and many species populations. I restrict myself to mention only some results from the very nice and interesting work from 1980 although more general results might hold and are shown in [3]. However I present W. Alt's results in an own separated section, since it is a little bit difficult to get this reference.

The time-dependent Keller-Segel system is included in the class of quasilinear parabolic cross-diffusion systems and the steady state problem belongs to the class of quasilinear elliptic systems. Important tools in the studies of elliptic and parabolic equations of second order are comparison and maximum principles to prove qualitative properties of the solution like boundedness or blow-up phenomena of the solution by constructing suitable super- and subsolutions for the considered problems. Also for existence results for elliptic and parabolic problems comparison principles have been used to apply Perron's method. Wolfgang Alt presents such comparison principles in [3] which also hold for coupled systems of the following general form:

$$B_0(\mathbf{u}(y))D_0\mathbf{u}(y) = \sum_{i,j=1}^n \frac{\partial}{\partial y_i} \left( a_{i,j}(y)A(\mathbf{u}(y))\frac{\partial}{\partial y_j}\mathbf{u}(y) \right) - \sum_{k=1}^l B_k(\mathbf{u}(y))\sum_{\mu=1}^m b_{k,\mu}(y)D_\mu\mathbf{u}(y)$$

$$(38) + C(y, \mathbf{u}(y), D\mathbf{u}(y)) + F(y, \mathbf{u}(y))$$

on a domain  $\Lambda$  in  $\mathbb{R}^d$  and for a function  $\mathbf{u} \in C^2(\Lambda, \mathbb{R}^M)$  resp. a distribution  $\mathbf{u} \in \mathcal{D}'(\Lambda, \mathbb{R}^M)$ , where

1. The notations

 $D_{\mu}\mathbf{u}(y) := X_{\mu}(y) \cdot \nabla \mathbf{u}(y)$  and  $Du(y) := (D_{1}\mathbf{u}(y), ..., D_{m}\mathbf{u}(y)) \in \mathbb{R}^{m,M}$ 

are used for vectorfields  $X_{\mu} = (\beta_1^{\mu}, ..., \beta_n^{\mu}) \in C^{1,1}_{loc}(\overline{\Lambda}, \mathbb{R}^M)$   $(\mu = 0, 1, ..., m)$  and  $y \in \overline{\Lambda}$ .

2. The  $a_{i,j}$  are defined as  $a_{i,j}(y) = \sum_{\nu,\mu=1}^{m} a_{\nu,\mu}(y)\beta_i^{\mu}(y)\beta_j^{\mu}(y)$  with continuous functions  $a_{\mu,\nu} = a_{nu,\mu}$  and  $(a_{\mu,\nu}) \ge 0$  on  $\overline{\Lambda}$ , and satisfy for all  $y \in \overline{\Lambda}$  and all  $\xi \in \mathbb{R}^m$  the inequality

$$\sum_{\nu,\mu=1}^{m} \xi_{\nu} \xi_{\mu} a_{\nu,\mu}(y) \ge \zeta(y) |\xi|^{2}$$

with a on  $\overline{\Lambda}$  lower semicontinuous positive function  $\zeta$ .

- 3. The functions  $b_{i,\mu}$  are continuous on  $\overline{\Lambda}$ .
- 4. The matrix-functions  $A, B_i : \mathbb{R}^M \to \mathbb{R}^{M \times M}$  are continuous and  $\det(A) \ge 0$  and  $B_0 \ge 0$  on  $\mathbb{R}^M$ .
- 5.  $F: \overline{\Lambda} \times \mathbb{R}^M \to \mathbb{R}^M$  and  $C: \overline{\Lambda} \times \mathbb{R}^M \times \mathbb{R}^{m,M} \to \mathbb{R}^M$  are measurable in  $y \in \Lambda$ , continuous in  $z \in \mathbb{R}^M$  and  $w \in \mathbb{R}^{m,M}$ , and uniformly bounded on  $\overline{\Lambda} \times \mathcal{K}$ , with a compact set  $\mathcal{K} \subset \mathbb{R}^M$ . For example the function C can be given by  $\mathbb{R}^M$ -valued bilinear forms  $C_{u,v}$  like:

$$C(y, z, w) = \sum_{\mu, \nu=1}^{m} C_{\mu, \nu}[w^{\mu}, w^{\nu}]$$

with  $w = (w^1, ..., w^m) \in \mathbf{I\!R}^{M, m}$ .

Although W. Alt's results hold for systems in this general setting I restrict myself to systems appropriate to model chemotaxis and present his results if possible in versions for those problems. From the applicational point of view one would like to know whether the considered model remains bounded for all times or not. Thus the existence of invariant sets for the system is an interesting topic worth studying. Alt presents such results in his Habilitation. Therefore the first result presented here is the following invariance theorem for parabolic systems (see [3, Satz 1.25, page 31 & 32]):

**Theorem 19 (Alt)** Let  $p \ge 0$  be in  $C_{loc}^{1,1}(\mathbb{R}^M)$  and let  $\mathcal{M}$  be defined as the set  $\mathcal{M} := \{z \in \mathbb{R}^M \mid p(z) = 0\} \neq \emptyset$ . Furthermore let us assume that **u** is a weak solution of the parabolic problem

$$\mathbf{u}_{t} = \sum_{j=1}^{N} \frac{\partial}{\partial x_{j}} \left( A(\mathbf{u}) \frac{\partial}{\partial x_{i}} \mathbf{u} \right) + F(\mathbf{u}), \quad in \ \Omega \times (0, \tau) \ (\Omega \subset \mathbf{I} \mathbf{R}^{N})$$
$$A(\mathbf{u}) \frac{\partial \mathbf{u}}{\partial n} = \psi(\mathbf{u}), \qquad on \ \partial \Omega_{N} \times [0, \tau]$$
$$\mathbf{u}(x, 0) = \mathbf{u}_{0}(x), \qquad on \ \partial \Omega_{D} \times [0, \tau],$$

where  $\partial \Omega_N$  and  $\partial \Omega_D$  denote disjunct subsets of  $\partial \Omega$ . For the boundary conditions we assume that there exists a continuous family of symmetric  $M \times M$ -matrices  $A^*(z) \ge 0, z \in \mathbb{R}^M$  and that there exist vector functions  $\theta_i : \partial \Omega_N \times \mathbb{R}^M \to \mathbb{R}$  such that

$$\psi_j(x,z) = A^*(z)\theta_j(x,z)$$
 and  $\theta_j^T(x,z) \cdot A^*(z) \cdot \theta_j(x,z) \leq const_{\mathcal{K}}$ 

for all z in a compact subset  $\mathcal{K}$  of  $\mathbb{R}^M$  and  $x \in \partial \Omega_N$ , where  $\psi_0 = \psi$  and  $\psi_j = \frac{\partial}{\partial z_j} \psi$ , for j = 1, ..., M. Furthermore we suppose that there exists a neighbourhood  $\mathcal{U}$  of  $\mathcal{M}$  such that

 $\nabla p \cdot \psi \leq 0, \ \nabla p \cdot F \leq 0 \text{ as well as } \nabla^2 p \cdot A \geq 0 \text{ holds on } U \setminus \mathcal{M}, \text{ if } \nabla^2 p \text{ exists.}$ 

*Then*  $\mathbf{u}(\cdot, 0) \subset \mathcal{M}$  *implies*  $\mathbf{u}(\cdot, t) \subset \mathcal{M}$  *for all*  $t \in [0, \tau]$ .

However not only the question whether the solution remains in a certain set for all times if the initial data belongs to this set is an interesting one. As demonstrated in the previous section the existence of Lyapunov functionals is a helpful tool in the studies of the time asymptotic as  $t \to \infty$ . In [3] W. Alt presents results for the existence of Lyapunov functionals which are different from the results of Theorem 17 of this paper. To be precise W. Alt proved the following Corollary:

**Corollary 1 (Alt)** Assume that the assumptions of Theorem 19 are fulfilled. The Dini-derivative of the functional

$$E(t) := \int_{\Omega} p(\mathbf{u}(t, x)) dx, \ t \in [0, \tau]$$

satisfies for all weak solutions which have values in U the inequality

$$\frac{d}{dt}E(t) \leq -\int_{\Omega} \sum_{j=1}^{N} \left(\frac{\partial}{\partial x_{j}}\mathbf{u}\right) \nabla^{2} p(\mathbf{u}) A(\mathbf{u}) \left(\frac{\partial}{\partial x_{j}}\mathbf{u}\right) dx + \int_{\Omega} \nabla p(\mathbf{u}) \cdot F(\mathbf{u}) dx + \int_{\partial\Omega} \nabla p(\mathbf{u}) \cdot \psi(\mathbf{u}) dS$$

almost everywhere in  $[0, \tau]$ . If for all  $z \in U \setminus M$  either  $\nabla p \cdot \psi < 0$  or  $\nabla p \cdot F < 0$  holds, then E is a Lyapunov functional for M, i.e.  $\frac{d}{dt}E(t) < 0$  as long as E(t) > 0. If for all  $z \in U \setminus M$  either  $\nabla p \cdot \psi < 0$  or  $\nabla p \cdot F < 0$  or  $\nabla^2 p \cdot A < 0$  holds, then E is a Lyapunov functional for  $M \cup N_0$ , where  $N_0$  contains all constants  $z_0 \in \mathbb{R}^M$ , that are zeros of  $\nabla p \cdot \psi$  and  $\nabla p \cdot F$ .

Now, let me demonstrate this results by applying Theorem 19 to an example (see [3, Beispiel 1.41, page 38]):

Consider the weak solution of the taxis-system

$$(39) \begin{cases} u_t = \nabla(k_1(u)(\nabla u - k_2(u, v)\nabla v)) + f(u, v), & \text{in } \Omega \times \{t \ge 0\} \\ v_t = k_c \Delta v + g(u, v), & \text{in } \Omega \times \{t \ge 0\} \\ 0 = k_1(u)\frac{\partial u}{\partial n} - h(u, v) & \text{on } \partial\Omega \times \{t \ge 0\} \\ 0 = \frac{\partial v}{\partial n}, & \text{on } \partial\Omega \times \{t \ge 0\} \end{cases}$$

where all coefficient functions are continuous and  $k_1(u), k_c \ge 0, k_1(u) = 0$  for  $u \ge \overline{u}$ . Furthermore let  $f(u, v), h(u, v) \le 0$  for  $u \ge \overline{u}$  and all  $v \in \mathbb{R}$  and on every compact subset of  $\mathbb{R}^2$  the inequalities

$$|h(u,\cdot)|^2, |\frac{\partial}{\partial u}h(u,\cdot)|^2, |\frac{\partial}{\partial v}h(u,\cdot)|^2 \le k_1(u)$$

hold and assume that the initial data  $(u_0, v_0)$  satisfy  $u_0 \leq \overline{u}$ .

We see that this system satisfies the conditions of Theorem 19 with

$$\mathcal{M} = \{(u, v) \in \mathbb{R}^2 \mid u \leq \overline{u}\}, \ p(u, v) = \frac{1}{2} \left( \max\{0, u - \overline{u}\} \right)^2$$

and

$$A(u,v) = \begin{pmatrix} k_1(u) & -k_1(u)k_2(u,v) \\ 0 & k_c \end{pmatrix}.$$

Now we have to check the conditions for the boundary data  $\psi = (h, 0)^T$ . We set

$$A^* = \begin{pmatrix} k_1 & 0\\ 0 & \frac{1}{2}(k_c + k_1 k_2^2) \end{pmatrix}.$$

Thus we have  $\psi = A^*\theta$  with  $\theta = (h/k_1, 0)^T$ . We see that  $\theta^T A^*\theta$  can be bounded uniformly and that the analoguous statements also hold for the partial derivatives  $\frac{\partial}{\partial u}\psi$  and  $\frac{\partial}{\partial y}\psi$ . Thus we conclude with Theorem 19 the following:

If the initial data  $(u_0, v_0)$  satisfy  $u_0 \le \overline{u}$ , then the solution of system (39) satisfies  $u(\cdot, t) \le \overline{u}$  for all  $t \ge 0$ .

In contrast to this result for weak solutions the next Theorem [3, Satz 2.40] is a strong (local) comparison principle for classical solutions of system (38). Before citing this result we have to introduce "sets of comparison" and "comparison surfaces".

**Definition 3** For i = 1, ..., I and a domain  $\mathcal{V} \subset \mathbb{R}^M$  let the functions  $\varrho_i$  belong to the class  $C_{loc}^{1,1}(\mathcal{V}, \mathbb{R}^M)$ , where  $\varrho_i$  is piecewise of class  $C_{loc}^{2,1}(\mathcal{V}, \mathbb{R}^M)$  and the functions  $\varsigma_i$  belong to the class  $C_{loc}^{1,1}(\overline{\mathcal{V}}, \mathbb{R})$ . Then we define for each  $y \in \overline{\mathcal{V}}$  the sets of comparison

 $\mathcal{M}_{\mathcal{Y}} := \{ z \in \mathcal{V} \mid \varrho_i(z) \le \varsigma_i(z), \ i = 1, ..., I \}$ 

and for i = 1, ..., I the  $\varrho_i$ -boundary

 $\partial_i \mathcal{M}_y := \{ z \in \partial \mathcal{M}_y \subset \mathcal{V} \mid \varrho_i(z) = \varsigma_i(z) \}.$ 

**Definition 4** Let  $(\mathcal{V}_y)_{y \in \Lambda}$  denote a continuous family of sets  $\mathcal{V}_y \subset \mathcal{V} \subset \mathbb{R}^M$ , with  $\mathcal{V}$  as in the previous definition. Furthermore let there be given functions  $\varrho \in C^{1,1}_{loc}(\overline{\mathcal{V}})$  and  $\varsigma \in C^{1,1}_{loc}(\Lambda)$  such that for each compact set  $K \subset \mathbb{R}^M$  and for each relatively compact set  $\Lambda^* \subset \Lambda$  there exist positive constants c and  $\delta$  and a continuous family of projections

$$\pi_{y}: \mathcal{M}_{y}^{\delta} := \mathcal{V}_{y} \cap K \cap \{\varsigma(y) - \delta \le \varrho \le \varsigma(y)\} \to \mathcal{V}_{y} \cap \{\varrho = \varsigma(y)\}$$

with

$$|\pi_{y}(z) - z| \leq c|\varrho(z) - \varsigma(y)|$$
 and  $\nabla \varrho(z) \neq 0$ 

for all  $y \in \Lambda^*$  and  $z \in \mathcal{M}_{v}^{\delta}$ . Furthermore let there be a finite number of sets  $\mathcal{V}^{\kappa}$  with

$$\mathcal{V} = \bigcup_{\kappa=1}^{k_0} \mathcal{V}^{\kappa} \text{such that } \varrho | \overline{\mathcal{V}^{\kappa}} \in C^{2,1}_{loc}(\overline{\mathcal{V}^{\kappa}}) \text{ and}$$
$$\pi_{\nu}(\overline{\mathcal{V}^{\kappa}} \cap \mathcal{V}_{\nu}) \subset \overline{\mathcal{V}^{\kappa}}$$

for all  $\kappa = 1, ..., k_0$  and  $y \in \Lambda^*$  is satisfied. Then we will call the surfaces  $\mathcal{V}_y \cap \{\varrho = \varsigma(y)\}$  surfaces of comparison.

Now let us turn to the strong comparison result for the equation

(40) 
$$B_{0}(\mathbf{u}(y))D_{0}\mathbf{u}(y) = A(\mathbf{u}(y))\sum_{\nu,\mu=1}^{m} a_{\nu,\mu}(y)D_{\nu}D_{\mu}\mathbf{u}(y) - \sum_{i=1}^{l} B_{i}(\mathbf{u}(y))\sum_{i,\mu=1}^{m} b_{i,\mu}(y)D_{\mu}\mathbf{u}(y) + \sum_{\nu,\mu=1}^{m} C_{\nu,\mu}(y,\mathbf{u}(y))(D_{\mu}\mathbf{u}(y) \cdot D_{\nu}\mathbf{u}(y)) + F(y,\mathbf{u}(y))$$

where we additionally assume that A,  $B_i$ , F and  $C_{\nu,\mu}$  are locally Lipschitz continuous in **u** and uniformly continuous in  $y \in \overline{\Lambda}$ , and that the functions  $b_{i,\mu}$  are bounded on  $\overline{\Lambda}$ :

**Theorem 20 (Alt)** Let us assume that  $(\mathcal{M}_y)_{y\in}\overline{\Lambda}$  is a family of sets of comparison. Additionally we assume that for each  $z \in \mathcal{V}$ , for which  $y \in \overline{\Lambda}$  and  $i \in \{1, ..., I\}$  exist such that  $z \in \partial_i \mathcal{M}_y$ , the following properties are satisfied:

$$\nabla \varrho_i(z) \neq 0, \ \nabla \varrho_i(z) \notin \sum_{s \neq i, z \in \partial_s \mathcal{M}_y} \mathbf{I} \mathbf{R} \cdot \nabla \varphi_s(z)$$

and the points of discontinuity of  $\nabla^2 \varrho_i$  lie on a finite system of smooth surfaces in  $\mathcal{V}$ , which intersect the surface  $\{z \in \mathcal{V} \mid \varrho_i(z) = \varsigma_i(z)\}$  in each such  $z \in \partial_i \mathcal{M}_y$  transversally. Let us suppose that for each pair  $(\varrho_i, \varsigma_i)$  and the families of sets  $(\mathcal{V}_y^i)_{y \in \Lambda}$  with

$$\mathcal{V}_y^l := \{z \in \mathcal{V} \mid \varrho_l(z) \leq \varsigma_l(z), \ l \in \{1, ..., I\} \setminus \{i\}\}$$

there exist continuous functions  $\zeta_i$  and  $\iota_i$  on  $\Lambda \times \mathcal{V}$  such that the following three properties are satisfied on  $\partial_i \mathcal{M}_y$ ,  $y \in \Lambda$ :

1. There are continuous functions  $\lambda^i$ ,  $\lambda^i_0$ , ...,  $\lambda^i_s$  on  $\mathcal{V}$  such that for all  $z \in \mathcal{V}_y$  with  $\varrho_i(z) = \varsigma_i(y)$  and  $y \in \Lambda$  such that

(41) 
$$\nabla \varrho_i(z) \cdot A(z) = \lambda^i(z) \nabla \varrho_i(z), \ \lambda^i(z) > 0$$

- (42)  $\nabla \varrho_i(z) \cdot B_0(z) = \lambda_0^i(z) \nabla \varrho_i(z), \ \lambda_0^i(z) > 0$
- (43)  $\nabla \varrho_i(z) \cdot B_l(z) = \lambda_l^i(z) \nabla \varrho_i(z) + p_l(z), \ \lambda_l^i(z) \in \mathbb{R}$
- (44) and  $p_l(z) \in [\nabla \varrho_i(z)]^{\perp} \subset \mathbf{I}\!\mathbf{R}^M, l = 1, ..., s.$
- 2. Let  $C(y,z) \subset \mathbb{R}^{m,M}$ ,  $(z \in V)$  denote for every  $y \in \Lambda$  a (uniform in  $y \in \Lambda^*$ ) locally Lipschitz continuous, given family of sets. The matrix  $W(y) \in \mathbb{R}^{m,m}$  denotes the positive definite root of the symmetric matrix  $(a_{\nu,\mu}(y))$  and  $\wp_i$  denotes for each  $i \in \{1,..,I\}$  a continuous vector field on V such that  $\wp_i(z) \cdot \nabla \varrho_i(z) = 1$  for all  $z \in V$ . Furthermore let us assume that for all  $y \in \Lambda$ ,  $z \in V_y$  with  $\varrho_i(z) = \varsigma_i(y)$  and  $w \in P_z(W(y) \cdot C(x,z)) := ((W(y) \cdot C(x,z)) - ((W(y) \cdot C(x,z)) \cdot \nabla \varrho_i(z)) \wp_i(z))$

(45) 
$$\lambda^{i}(z) \sum_{k=1}^{m} \min_{\kappa=1,\dots,k_{0} \ ; \ z \in} \overline{\mathcal{V}^{\kappa}} \{ w_{k} \cdot \nabla^{2} \varrho_{i}(z) \cdot w_{k} \} + \sum_{j=1}^{l} \sum_{k=1}^{m} \left( \sum_{\mu=1}^{m} b_{j,\mu}(y) \mathcal{W}_{\mu,k}^{-1}(y) \right) w_{j}(z) \cdot w_{k}$$
$$- \sum_{\nu,\mu,k,l=1}^{m} \mathcal{W}_{\mu,k}^{-1}(y) \mathcal{W}_{\nu,l}^{-1}(y) \nabla \varrho_{i}(z) \cdot C_{\mu,\nu}(y,z) |w_{k},w_{l}] \ge \zeta_{i}(x,z) |w|^{2} - \iota_{i}(y,z)$$

3. Let  $S(y_0, \Lambda)$  denote the set of all points  $y \in \Lambda$  such that there exists a continuous, piecewise  $C^{1,1}$  curve  $\gamma$  with  $\gamma(0) = y_0$  and  $\gamma(t) \in \Lambda$  for all t > 0, which is the inte-

gral-curve to one of the vector fields  $-X_0, \pm X_1, ..., \pm X_m$ . The generalized Hessian matrix  $H^+_{\mu,\nu}\varsigma_i(x)$  of the second derivatives  $(D_{\mu}D_{\nu}\varsigma_i)$  of the function  $\varsigma_i$  is then defined as follows:

Denote for all  $\xi \in \mathbb{R}^m$  by  $\gamma_{\xi}$  the solution curve of the vector field  $\sum_{\mu=1}^m \xi_{\mu} X_{\mu}$  with  $\gamma_{\xi}(0) = y$ . Then we set

$$\sum_{\mu,\nu=1}^{m} \xi_{\mu}\xi_{\nu}H_{\mu,\nu}^{+}\varsigma_{i}(y) := d_{t}^{+}\frac{d}{dt}(\varsigma_{i}\circ\gamma_{\xi})\mid_{t=0}$$

with

$$d_t^+g(0):=\limsup_{t o 0}rac{g(t)-g(0)}{t}$$

for a Lipschitz continuous function g on  $\mathbb{R}$ .

Then, for all  $y \in \Lambda$  and  $z \in \mathcal{V}_y^i$  with  $\varrho_i(z) = \varsigma_i(y)$  the following comparison condition should hold:

$$\begin{split} \nabla \varrho_{i}(z) \cdot F(y,z) &+ \iota_{i}(y,z) \\ \leq \lambda_{0}^{i}(z) D_{0}\varsigma_{i}(y) - \lambda^{i}(y) \sum_{\mu,\nu=1}^{m} a_{\mu,\nu}(y) H_{\mu,\nu}^{+}\varsigma_{i}(y) + \sum_{j=1}^{l} \sum_{k=1}^{m} \lambda_{k}^{i}(z) b_{k,\mu}(y) D_{\mu}\varsigma_{i}(y) \\ &+ \sum_{\mu,\nu=1}^{m} \left( a_{\mu,\nu}\lambda^{i}(y) \min_{\substack{\kappa=1,\dots,k_{\sigma} \\ z \in \overline{\mathcal{V}^{\kappa}}}} (\wp_{i} \cdot \nabla^{2} \varrho_{i} \cdot \wp_{i})(z) + \nabla \varrho(z) \cdot C_{\mu,\nu}(y,z) [\wp_{i}(z), \wp_{i}(z)] \right) D_{\mu}\varsigma_{i}(y) D_{\nu}\varsigma_{i}(y) \\ &- \sum_{\mu,\nu=1}^{m} \left( \frac{c_{i}^{2}(y,z)}{\zeta_{i}(y,z)} \delta_{\mu,\nu} \right) D_{\mu}\varsigma_{i}(y) D_{\nu}\varsigma_{i}(y) \end{split}$$

*Here*  $c_i(y, z)$  *defines the function* 

(46) 
$$c_{i}(y,z) = \sup_{w \in P_{z}(W(y) \cdot C(x,z))} \left\{ \frac{1}{|w|} \sum_{k,\nu=1}^{m} \max_{k \; ; \; z \in \overline{\mathcal{V}^{k}}} (W_{k,\mu}(y)\lambda^{i}(y)\wp_{i}(y)\nabla^{2}\varrho_{i}(z)w_{k}) - \frac{1}{2} \sum_{\mu=1}^{m} W_{\mu,k}^{-1}(y)\nabla\varrho(z) \cdot (C_{\mu,\nu}(y,z)[w_{k},\wp_{i}(y)] + C_{\mu,\nu}(y,z)[\wp_{i}(z),w_{k}]) \right\}^{1/2}.$$

The case  $\zeta_i = 0$  and  $c_i > 0$  is allowed, if  $D_{\varsigma_i}$  can be chosen identically equal to zero.

Then for every solution  $\mathbf{u} \in C^2(\Lambda, \mathbb{R}^M)$  of (40) with  $D\mathbf{u}(y) \in C(y, \mathbf{u}(y))$  and  $\mathbf{u}(y) \in \mathcal{M}_y$  for all  $y \in \Lambda$  we have the following statements:

- 1. If there is a  $i \in \{1, ..., I\}$  and a  $y_i \in \Lambda$  with  $\mathbf{u}(y_i) \in \partial_i \mathcal{M}_{y_i}$  then  $\mathbf{u}(t) \in \partial_i \mathcal{M}_t$  for all  $t \in \overline{S(y_i, \Lambda)}$ .
- 2. Let  $J \subset \{1, ..., I\}$ , such that for every  $i \in J$  there exists a  $y_i \in \Lambda$  with  $\mathbf{u}(y_i) \in \partial_i \mathcal{M}_{y_i}$ , then  $\mathbf{u}(t) \in \bigcap_{i \in I} \partial_i \mathcal{M}_t$  for all  $t \in \bigcap_{i \in I} \overline{\mathcal{S}(y_i, \Lambda)}$ .

Theorem 20 is very general and technical. However it allows together with a reformulation of Hopf's maximum principle for systems (see [3, Lemma 3.2, page 63]) to find additional hypotheses under which the solution of certain boundary value problems remain in the interior of the sets of comparison. Furthermore it is possible to prove a general invariance theorem for parabolic systems. Here we only mention an application of these theorems to autonomous parabolic Neumann boundary value problems. Therefore let us once again consider the following problem:

$$\mathbf{u}_{t} = \sum_{j=1}^{N} \frac{\partial}{\partial x_{j}} \left( A(\mathbf{u}) \frac{\partial}{\partial x_{i}} \mathbf{u} \right) + F(\mathbf{u}), \quad \text{in } \Omega \times (0, \tau) \; (\Omega \subset \mathbf{I} \mathbf{R}^{N}$$
$$A(\mathbf{u}) \frac{\partial \mathbf{u}}{\partial n} = \psi(\mathbf{u}), \quad \text{on } \partial \Omega_{N} \times [0, \tau]$$

where we assume that  $F, \psi \in C^{0,1}_{loc}(\mathcal{V}, \mathbb{R}^M)$  and  $A \in C^{1,1}_{loc}(\mathcal{V}, \mathbb{R}^{M,M})$  with  $det(A) \ge 0$ .

**Theorem 21 (Alt)** Let  $\varrho_i \in C^{1,1}_{loc}(\mathcal{V}, \mathbb{R})$ ,  $\mathcal{V} \subset \mathbb{R}^M$  (i = 1, ..., I) denote given functions and let  $Q \subset \mathbb{R}^I$  be also given. For  $q \in Q$  the set  $\mathcal{M}_q$  is defined as

 $\mathcal{M}_q := \{ z \in \mathcal{V} \mid \varrho_i(z) \le q_i, \ i = 1, ..., I \}.$ 

Let  $q^0 \in Q$  be given, such that the initial condition  $u(0, x) \in \mathcal{M}_{q^0}$  is satisfied for all  $x \in \overline{\Omega} \times \{t = 0\}$ . Furthermore we assume that there exists a solution  $\omega \in C^1([0, \tau_0], \mathbb{R}^I)$  of the system of differential inequalities

$$\frac{d}{dt}\omega_i(t) \ge \sup_{z \in \partial_i \mathcal{M}_q} \{\nabla \varrho_i(z) \cdot F(z)\}, \ i = 1, .., I, \ 0 \le t \le \tau_0$$

such that  $\omega_i(0) \ge q_i^0$ , i = 1, ..., I and a positive, continuous function  $\zeta_0$  on  $[0, \tau_0]$ , such that the following conditions are satisfied for

$$Q := \{q \in \mathbb{R}^I \mid \exists t \in [0, \tau_0] \text{ with } \omega_i(t) \le q_i \le \omega_i(t) + \zeta_0(t) \text{ for all } i = 1, ..., I\}:$$

 $\nabla \varrho_i(z) \neq 0$  holds for all  $z \in \mathbf{R}^M$  with  $a \ q \in Q$  and  $z \in \partial_i \mathcal{M}_q$ . Furthermore  $\nabla \varrho_i(z) \notin \sum_{s \neq i, z \in \partial_s \mathcal{M}_q} \mathbf{R} \cdot \nabla \varphi_s(z)$  holds on the "edges" of  $\partial \mathcal{M}_q$  and  $\varrho$  is piecewise  $C^{2,1}$ . For all  $z \in \partial_s \mathcal{M}_q$  with  $a \in Q$  and  $i \in \{1, \dots\}$  we have

all  $z \in \partial_i \mathcal{M}_q$  with  $q \in Q$  and  $i \in \{1, .., I\}$  we have:

1.  $\nabla \varrho_i(z)A(z) = \lambda_i(z)\nabla \varrho_i(z), \ \lambda_i(z) > 0$ 2.  $\nabla^2 \varrho_i(z)A(z) \ge 0 \text{ on } [\nabla \varrho(z)]^{\perp}, \text{ for all } \kappa = 1, ..., k_0 \text{ with } z \in \overline{\mathcal{V}^{\kappa}}.$ 3.  $\nabla \varrho_i(z) \cdot \psi(z) \le 0.$ 

Then  $u(t, x) \in \mathcal{M}_{\omega(t)}$  for all  $(t, x) \in \Omega \times [0, \tau_0]$ .

This Theorem allows us to make statements on the time asymptotic behaviour of some special cases of the solution of Keller-Segel type models. For example (see also [3, Beispiel 5.18, pages 109–111]) let us consider the system

(47) 
$$\begin{cases} u_t = \nabla \left( \frac{k_1}{1+u} (\nabla u - \chi u) \nabla v \right) + (\beta - \alpha u), & \text{in } \Omega \times \{t \ge 0\} \\ v_t = \Delta v - \delta v + \gamma u, & \text{in } \Omega \times \{t \ge 0\} \\ \frac{\partial u}{\partial n} = 0 = \frac{\partial v}{\partial n}, & \text{on } \partial \Omega \times \{t \ge 0\} \end{cases}$$

## where $k_1 > 1$ and $\chi, \beta, \alpha, \delta, \gamma$ are positive constants. We define $m_0 = k_1 - 1 > \frac{\beta}{\alpha}$ and

$$\varrho(u,v) = v - \frac{m_0}{k_1 \chi} \log\left(\frac{u}{m_0}\right) + \frac{1}{k_1 \chi} (u - m_0)$$

Then

$$\nabla \varrho(u,v) = \left( c \frac{1}{k_1 \chi} \left( 1 - \frac{m_0}{u} \right) 1 \right) \text{ and } \nabla \varrho(u,v) \cdot F(u,v) = \frac{1}{k_1 \chi} (\beta - \alpha u) (1 - \frac{m_0}{u}) + \gamma u - \delta v.$$

We now set

$$\varrho_1(u, v) = -v, \ \varrho_2(u, v) = \varrho(u, v), \ \varrho_3(u, v) = v + \frac{1}{k_1 \chi} \left( u - m_0 - m_0 \log \left( \frac{u}{m_0} \right) \right)$$

and

$$q \in \mathbb{R}^3$$
 with  $q_1 = -a^-$ ,  $q_2 = a^+ - \frac{1}{k_1 \chi} \left( m_0 \log \left( \frac{\beta}{\alpha m_0} \right) - \frac{\beta}{\alpha} + m_0 \right)$  and  $q_3 = a^+$ 

for some  $0 \le a^- < a^+$ . The set of comparison  $\mathcal{M}_q$  is then convex. Now let us additionally assume that

$$0 < \gamma < \frac{\alpha - \delta}{k_1 \chi} \left( 1 - \frac{m_0}{\beta} \alpha \right)$$

holds. Let (u, v) denote the solution of the system (47) and let  $a_0^+ \ge \gamma m_0/\delta$  be minimal and  $a_0^- > 0$  be maximal such that with  $q^0$  defined analogously as above  $u(0, x) \in \mathcal{M}_{q^0}$  is satisfied for all  $x \in \Omega$ . If we set

$$a^{+}(t) = a_{0}^{+}e^{-\delta t} + \frac{\gamma m_{0}}{\delta} (1 - e^{-\delta t}), \ t \ge 0$$

and choose  $a^-$  as the solution of the ODE

$$\frac{d}{dt}a^{-}(t) = \gamma \underline{u}(t) - \delta a^{-}(t), \ a^{-}(0) = a_{0}^{-}$$

where  $\underline{u}(t) < \beta/\alpha$  is the uniquely defined by the equation

$$m_0 \log\left(\frac{\alpha \underline{u}(t)}{\beta}\right) + \frac{\beta}{\alpha} - \underline{u}(t) = -k_1 \chi a^+(t)$$

then we see for the analogously defined q(t) that  $u(t, x) \in \mathcal{M}_{q(t)}$  for all  $(t, x) \in \Omega \times [0, \infty)$ . The function  $a^+(t)$  converges as  $t \to \infty$  monotone decreasing to  $a^+_{\infty} = \gamma m_0/\delta$ , and the function  $a^-(t)$  converges as  $t \to \infty$  monotone increasing to a  $a^-_{\infty} \leq \gamma \beta/\delta \alpha < a^+_{\infty}$  which is the unique solution of the equation

$$m_0 \log\left(\frac{\delta \alpha a^-(t)}{\gamma \beta}\right) - \left(\frac{\delta}{\alpha} - k_1 \chi\right) a^-(t) + \frac{\beta}{\alpha} + \frac{k_1 \chi m_0 \gamma}{\delta} = 0.$$

Thus we see that the set  $M_{q_{\infty}}$  with the corresponding  $q_{\infty}$  is a global attracting set for all positive solutions of (47).

Wolfgang Alt's Habilitation from 1980 contains many results more. However a complete presentation of his nice results would expand the present work too much.

Thus this short presentation of some of his results should be enough. I refer the interested reader to [3] for more results, more applications and more examples of the invariance principles derived by W. Alt.

## 8 Conclusions and personal comments

There are of course still many open problems in connection with the various approaches that have been studied for the Keller-Segel model.

This paper dealt solely with the parabolic model proposed by Keller and Segel for the aggregation phase of mobile species caused by chemotaxis. However there are also different approaches to chemotaxis and hence also numerous different models describing chemotaxis. First of all, one must always keep in mind that the model one uses is based either on a microscopic or a macroscopic approach and always depends on the species studied. Thus transport and hyperbolic models for chemotaxis have also been proposed. For example transport and hyperbolic models for chemotactical movement have been studied in [4, 5, 35, 38, 39, 56, 58, 60] and [114]. The connection between chemotaxis equations as the parabolic limit of velocity jump processes or transport models for chemotaxis has been studied in [4, 5, 22, 59, 68, 117] and [118]. I refer to [38] for surveys on different models for chemotactical movement and to [60] for a survey on the hyperbolic approach to chemotaxis.

Of course there are many publications presenting experimental data on chemotactic effects and the influences of changes in the motility or the chemotactic sensitivity of the given species (see for example [36, 37]). Models which take the chemotactical movement of n populations according to k sensitivity agents into account have been proposed in [3, Beispiel 2.47, page 58], [32, 77, 80] and in [156, 157] by G. Wolansky.

Wolansky has studied a generalization of the Keller-Segel model for n populations in the absence of conflicts. He showed that under certain assumptions a conflict free system of n populations admits a Lyapunov functional. Using this functional he investigates the existence of steady state solutions via variational methods. Furthermore, he investigates time-periodic solutions in a parameter range where a Lyapunov functional does not exist.

In [32] a chemotaxtis system for two populations in case of a "conflict of interests" (according to the expressions in [157]) is studied. The system studied in [32] considers the two populations A and B where population A is attracted by a substance P and repelled by substance Q, while population B behaves the other way around. The populations are assumed to move chemotactical positive in direction of a higher concentration of their attractant and chemotactical negative away from higher concentrations of their repellent. The evolution equation for substance P depends only on population B and Q itself. Using an appropriate scaling and some mathematical necessary simplifying assumptions, the authors show that the existence of stationary solutions depends on the solvability of a two-parameter-dependent nonlinear second order boundary value problem with exponential nonlinearities. Furthermore the authors prove the existence of nontrivial solutions in the para-

meter-space. The presented bifurcation analysis in [32] shows that the total rescaled mass of the two populations plays the role of a bifurcation parameter for the existence of nontrivial stationary solutions.

In [77, 80] a model for n-populations is studied that takes competition and chemotaxis in a chemostat into account. I refer the interested reader to these two references for more details on that particular model and the results achieved.

Beside those results for the chemotactical movement of n populations, also the effect of multiple attractant gradients on chemotactical movement has been studied in [144] and numerical solutions for the corresponding models have been calculated in [40, 41]. Numerical analysis for the Keller-Segel model has been performed, for example, by Gajewski and Zacharias using a chemotaxis version of the well-established TOSCA code for solving semiconductor problems and by Nakaguchi and Yagi studying the full discrete approximation of the Keller-Segel model by Galerkin Runge-Kutta methods [104, 105]. The transport chemotaxis model is dealt with in [21] and [41], for example.

The results available for systems related to the Keller-Segel equations such as the Othmer-Stevens model are so numerous that I mention only a few [81, 116, 145] and [159]. The well-posedness of the Othmer-Stevens model follows directly from the results by Rascle [121].

Of course, the functional forms appearing in the original Keller-Segel model can vary from species to species and some explicit models for the cAMP oscillation have been proposed (see [84] for the so called Martiel and Goldbeter model, [115] for a survey on the oscillatory cAMP signaling in Dictyostelium discoideum and [124] for the description of the role of cAMP in the development of Dictyostelium discoideum). The Keller-Segel model has also been used to describe different problems. For example, in [88] the Keller-Segel equations have been proposed for strip pattern formation in alligator embryos. Angiogenesis has also been proposed as another application of Keller-Segel type models (see for example [27] including references). The large number of applications and of possible functional forms results directly in a large number of models depending on the considered problem. In some particular papers this has also resulted in the addition of a third equation to the Keller-Segel model (4) or the rediscovery of a more complicated version of the Keller-Segel model. This results from attempts to describe more complicated pattern formations during the aggregation phase of mobile species such as the attempt to describe spiral waves during the aggregation (see, for example, [138, 148] and [149] for such extended models).

At the conclusion of this survey I would ask the reader to allow me a personal comment. The references given in this text are far from complete. I have tried to give the interested reader a brief summary of the latest developments in the Keller-Segel model. Thus, this article is intended as continuation of Evelyn Fox Keller's article "Assessing the Keller-Segel model: How has it fared" of 1980 [76]. It is left to the reader to decide whether I have succeeded.

Acknowledgement: I thank Prof. Dr. Wolfgang Alt for allowing me to include the results from his Habilitation in this summary of the Keller-Segel system.

## References

- [1] J. Adler, Chemotaxis in bacteria. Ann. Rev. Biochem. 44 (1975), pp. 341-356.
- [2] N. D. Alikakos, L<sup>p</sup> bounds of solutions of reaction-diffusion equations Comm. In Partial Differential Equations, 4 (1979), pp. 827–868.
- [3] W. Alt, Vergleichsätze für quasilineare elliptisch-parabolische Systems partieller Differentialgleichungen, Habilitation, Ruprecht-Karl-Universität Heidelberg, 1980.
- [4] W. Alt, Biased Random Walk models for chemotaxis and related diffusion approximations, J. Math Biol. 9 (1980), pp. 147–177.
- [5] W. Alt, Singular perturbation of differential integral equations describing biased random walks. J. Reine Angew. Math. 322 (1981), pp. 15 – 41.
- [6] H. Amann, Dynamic theory of quasilinear parabolic equations II. Reaction-Diffusion Systems, Differential and Integral Eqns. 3 (1990), pp. 13–75.
- [7] H. Amann, Nonhomogeneous linear and quasilinear elliptic and parabolic boundary value problems, Function Spaces, Differential Operators and Nonlinear Analysis (Eds. Schmeisser, Triebel), Teubner Texte zur Mathematik 133 (1993), pp. 9–126.
- [8] F. Berezovskaya and G. Karev, Bifurcation of travelling waves in population taxis models. Physics-Uspekhi, 42 (1999) 917–929.
- [9] F. Berezovskaya and G. Karev, Parametric portraits of travelling waves of population models with polynomial growth and autotaxis rates. Spatial heterogeneity in ecological models (Alcalß de Henares, 1998). Nonlinear Anal. Real World Appl. 1 (2000) 123–136.
- [10] F. Berezovskaya and G. Karev, Polynomial models of populations with autotaxis: "traveling wave" solutions. (Russian). Mat. Model. 12 (2000) 78–93.
- [11] P. Biler, Local and global solvability of some parabolic system modelling chemotaxis, Adv. Math. Sci. Appl. 8 (1998), pp. 715–743.
- [12] P. Biler, Global solutions to some parabolic-elliptic systems of chemotaxis, Adv. Math. Sci. Appl. 9 (1999), pp. 347–359.
- [13] A. Bonami, D. Hilhorst, E. Logak, and M. Mimura, A free boundary problem arising in a chemotaxis model, Free boundary problems, theory and applications (Zakopane, 1995), Pitman Res. Notes Math. Ser. 363, Longman, Harlow (1996), pp. 368–373.
- [14] A. Bonami, D. Hilhorst, E. Logak, and M. Mimura, Singular limit of a chemotaxis-growth model, Adv. Differential Equations 6 (2001), pp. 1173–1218.
- [15] J. T. Bonner, *The cellular slime molds*, Princeton University Press, Princeton, New Jersey, second edition, 1967.
- [16] A. Boy, Analysis for a system of coupled reaction-diffusion parabolic equations arising in biology. Computers Math. Applic 32 (1996), pp. 15–21.
- [17] M. P. Brenner, L. S. Levitov and E. O. Budrene, *Physical mechanisms for chemotactic pattern formation by bacteria*. Biophys. J. 74 (1998), pp. 1677–1693.
- [18] M. P. Brenner, P. Constantin, L. P. Kadanoff, A. Schenkel and S. C. Venkataramani, Attraction, Diffusion and Collapse, Nonlinearity 12 (1999), pp. 1071–1098.
- [19] M. P. Brenner, P. Constantin, L. P. Kadanoff, A. Schenkel and S. C. Venkataramani, Blowup in the Chemotaxis equation, Technical Report University of Chicago.
- [20] H. Brézis and F. Merle, Uniform estimates and blow-up behaviour for solutions of  $-\Delta u = V(x)e^{u}$  in two dimensions, Comm. P.D.E. 16 (1991), pp. 1223–1253.
- [21] B. J. Brosilow, R. M. Ford, S. Sarman and P. T. Cummings, Numerical solution of transport equations for bacterial chemotaxis: Effect of discretization of directional motion, SIAM J. Appl. Math. 56 (1996), pp.1639–1663.
- [22] F. A. C. C. Chalub, P. Markowich, B. Perthame and C. Schmeiser, Kinetic models for chemotaxis and their drift diffusion limit. Preprint 2002.
- [23] S.-Y.A. Chang and P. Yang, Conformal deformation of metrics on S<sup>2</sup>, J. Differential Geometry, 27 (1988), pp. 259–296
- [24] S. Childress and J.K. Percus, Nonlinear aspects of chemotaxis, Math. Biosc. 56 (1981), pp. 217-237.

Übersichtsartikel

- [25] S. Childress, Chemotactic collapse in two dimensions, Lecture Notes in Biomathematics, Springer-Verlag, volume 55, 1984, pp. 61–66.
- [26] M. H. Cohen and A. Robertson, Wave propagation in the early stages of aggregation of cellular slime molds. J. theor. Biol. 31 (1971), 101–118.
- [27] L. Corrias, B. Perthame and H. Zaag, A model motivated by angiogenesis, C. R. Acad. Sci. Paris, Ser. I 336 (2003), pp. 141–146.
- [28] B. Davis, Reinforced random walks, Probab. theory Related Fields 84 (1990), pp. 203-229.
- [29] F. Dkhil, On the analysis of reaction-diffusion-advection systems arising in chemical and biomathematical models, PhD-thesis, University of Cergy-Pontoise, France, 2002.
- [30] Y. Ebihara, Y. Furusho and T. Nagai, Singular solutions of traveling waves in a chemotactic model., Bull. Kyushu Inst. Tech. Math. Natur. Sci. 39 (1992) 29–38.
- [31] Evans, L. C. 1998 Partial Differential Equations, American Mathematical Society, Graduate Studies in Mathematics 19, Providence, Rhode Island.
- [32] A. Fasano, A. Mancini and M. Primicerio, Equilibrium of two populations subjected to chemotaxis, Preprint 2002.
- [33] D. L. Feltham and M. A. J. Chaplain, Travelling waves in a model of species migration, Appl. Math. Lett. 13 (2000), pp. 67–73.
- [34] P. C. Fife and J. B. McLeod, The approach of solutions of nonlinear diffusion equations to traveling front solutions, Arch. Rat. Mech. Analysis 65 (1977), pp. 335–362.
- [35] R. M. Ford and D. A. Lauffenburger, Analysis of chemotactic bacterial distributions in population migration assays using mathematical model applicable to steep or shallow attractant gradients, Bull. Math. Biol. 53 (1991), pp. 721–749.
- [36] R. M. Ford, B. R. Phillips, J. A. Quinn and D. A. Lauffenburger, Measurement of bacterial random motility and chemotaxis coefficients: I. Stopped-flow diffusion chamber assay, Biotech. and Bioengin. 37 (1991), pp. 647–660.
- [37] R. M. Ford and D. A. Lauffenburger, Measurement of bacterial random motility and chemotaxis coefficients: II. Application of Single-cell-based mathematical model, Biotech. and Bioengin. 37 (1991), pp. 661–672.
- [38] R. M. Ford, Mathematical modeling and quantitative characterization of bacterial motility and chemotaxis., Modelin the Metabolic and Physiology Activities of Microorganisms: Chapter 7, John Wiley & sons, Inc. 1992.
- [39] R. M. Ford and P. T. Cummings, On the relationship between cell balance equations for chemotactic cell populations, SIAM J. Appl. Math. 52 (1992), pp. 1426–1441.
- [40] P. D. Frymier, R. M. Ford and P. T. Cummings, Cellular dynamics simulations of bacterial chemotaxis, Chem. Engin. Sci. 48 (1993), pp. 687–699.
- [41] P. D. Frymier, R. M. Ford and P. T. Cummings, Analysis of bacterial migration: I. Numerical solution of balance equation, AIChE J. 40 (1994), pp. 704–715.
- [42] M. Funaki, M. Mimura and T. Tsujikawa, *Traveling front solutions arising in a chemotaxis-growth model*, RIMS Kokyuroku 1135 (2000), pp. 52–76.
- [43] H. Gajewski, W. Jäger and A. Koshelev, About loss of regularity and "blow up" of solutions for quasilinear parabolic systems, Preprint No. 70, Weierstraß-Institut für Angewandte Analysis und Stochastik, Berlin (1993).
- [44] H. Gajewski and K. Zacharias, Global behavior of a reaction-diffusion system modelling chemotaxis, Math. Nachr. 195 (1998), pp. 77-114.
- [45] H. Gajewski and K. Zacharias, On a reaction-diffusion system modelling chemotaxis, Proceedings of the Equadiff 99, (eds. B. Fiedler, K. Gröger, J. Sprekels), World Scientific Publishing Co, Singapore, 2000, pp. 1098–1103.
- [46] J. Grolle (verantwortlicher Redakteur f
  ür Biologie), Wabernder Klecks, DER SPIEGEL 36 (1998), p. 212.
- [47] C. Gui, J. Wei and M. Winter, Multiple boundary peak solutions for some singulary perturbed Neumann problems, Ann. Inst. H. Poincaré Anal. Non Linéaire 17 (2000), pp. 47–82.
- [48] G. Harada, T. Nagai, T. Senba and T. Suzuki, Concentration lemma, Brezis-Merle type inequality, and a parabolic system of chemotaxis, Adv. Differential Equations 6 (2001), pp. 1255–1280.

- [49] M. Henry, D. Hilhorst and R. Schätzle, Convergence to a viscosity solution for an advection-reaction-diffusion equation arising from a chemotaxis-growth model. Hiroshima Math. J. 29 (1999), pp. 591–630.
- [50] M. A. Herrero and J. J. L. Velázquez, Singularity patterns in a chemotaxis model, Math. Ann. 36 (1996), pp. 583-623.
- [51] M. A. Herrero and J. J. L. Velázquez, Chemotactic collapse for the Keller-Segel model, J. Math. Biol. 35 (1996), pp. 583-623.
- [52] M. A. Herrero and J. J. L. Velázquez, A blow-up mechanism for a chemotaxis model, Ann. Scuola Normale Superiore 24 (1997), pp. 633–683.
- [53] M. A. Herrero, E. Medina and J. J. L. Velázquez, Finite-time aggregation into a single point in a reaction diffusion system, Nonlinearity 10 (1997), pp. 1739–1754.
- [54] M. A. Herrero, E. Medina and J. J. L. Velázquez, Self-similar blow-up for a reaction diffusion system, Journal of Computational and Applied Mathematics 97 (1998), pp. 99–119.
- [55] M. A. Herrero, Asymptotic properties of reaction-diffusion systems modelling chemotaxis, Applied and industrial mathematics, Venice-2, 1998, Kluwer Acad. Publ., Dordrecht 2000, pp. 89–108.
- [56] T. Hillen and A. Stevens, *Hyperbolic models for chemotaxis in 1-D*, Nonlinear Anal., Real World Appl. 1 (2000), pp. 409–433.
- [57] T. Hillen and K. J. Painter, Global Existence far a Parabolic Chemotaxis Model with Prevention of Overcrowding, Adv. in Appl. Math. 26 (2001), pp. 280–301.
- [58] T. Hillen, C. Rohde and F. Lutscher Existence of weak solutions for a hyperbolic model of chemosensitive movement, J. Math. Anal. Appl. 260 (2001), pp. 173–199.
- [59] T. Hillen and H. G. Othmer, The diffusion limit of transport equations derived from velocityjump processes, SIAM J. Appl. Math. 61 (2000), pp. 751–775.
- [60] T. Hillen, Hyperbolic models for chemosensitive movement, Math. Models Methods Appl. Sci. 12 (2002), pp. 1–28.
- [61] D. Horstmann, Some blowup results for the Keller-Segel model, Proceedings of the Equadiff 99, (eds. B. Fiedler, K. Gröger, J. Sprekels), World Scientific Publishing Co, Singapore 2000, pp. 1104–1106.
- [62] D. Horstmann, Lyapunov functions and L<sup>p</sup>-estimates for a class of reaction-diffusion systems, Colloquium Mathematicum 87 (2001), pp. 113–127.
- [63] D. Horstmann, The nonsymmetric case of the Keller-Segel model in chemotaxis: some recent results, Nonlinear Differential Equations and Applications (NoDEA) 8 (2001), pp. 399–423.
- [64] D. Horstmann and G. Wang, *Blow-up in a chemotaxis model without symmetry assumptions*, European Journal of Applied Mathematics **12** (2001), pp. 159–177.
- [65] D. Horstmann, On the existence of radially symmetric blow-up solutions for the Keller-Segel model, Journal of Mathematical Biology 44 (2002), pp. 463–478.
- [66] D. Horstmann and A. Stevens, A constructive approach to traveling waves in chemotaxis, Preprint no.78: 2001 Max-Planck-Institute for Mathematics in the Sciences, Leipzig, submitted.
- [67] D. Horstmann, On some pattern forming effects in chemotaxis, to appear in the Proceedings of the 5th European Conference of the European Society for Mathematical and Theoretical Biology (ECMTB 2002).
- [67a]D. Horstmann, From 1970 until present: The Keller-Segel model in chemotaxis and its consequences II. Jahresber. Dtsch. Math.-Ver. (to appear).
- [68] H. J. Hwang, K. Kang and A. Stevens, *Drift-diffsion limits of kinetic models for chemotaxis: a generalization* Preprint no.19: 2003 Max-Planck-Institute for Mathematics in the Sciences, Leipzig, submitted.
- [69] W. Jäger and S. Luckhaus, On explosions of solutions to a system of partial differential equations modelling chemotaxis, Trans. Am. Math. Soc. 329 (1992), pp. 817–824.
- [70] Y. Kabeya and W.-M. Ni, Stationary Keller-Segel model with the linear sensitivity, RIMS Kokyuroku 1025 (1998), pp. 44–65.
- [71] E. F. Keller and L. A. Segel, *Initiation of slime mold aggregation viewed as an instability*, J. Theor. Biol. 26 (1970), pp. 399–415.
- [72] E. F. Keller and L. A. Segel, *Model for chemotaxis*, J. Theor. Biol. **30** (1971), pp. 225–234.

Übersichtsartikel

- [73] E. F. Keller and L. A. Segel, Traveling bands of chemotactic bacteria: A Theoretical Analysis, J. theor. Biol. 30 (1971), pp. 235–248.
- [74] E. F. Keller and G. M. Odell, Necessary and sufficient conditions for chemotactic bands Math. Biosci. 27 (1975), pp. 309–317.
- [75] E. F. Keller and G. M. Odell, Traveling bands of chemotactic bacteria revisited, J. Theor. Biol. 56 (1976), pp. 243–247.
- [76] E. F. Keller, Assessing the Keller-Segel modell: How has it fared?, Editors W. Jäger, H. Rost und P. Tautu, Lecture Notes in Biomathematics 38, Springer-Verlag, Berlin, Heidelberg, New York 1980, pp. 379–387.
- [77] H. J. Kuiper, A priori bounds and global existence for a strongly coupled quasilinear parabolic system modeling chemotaxis, Electron. J. Differ. Equ. 2001 (2001), pp. 1–18, electronic only.
- [78] O. A. Ladyzenskaya and N. N. Ural'ceva, *Linear and quasilinear elliptic equations*, Academic Press, New York and London, 1968.
- [79] D. Lauffenburger, C. R. Kennedy and R. Aris Traveling bands of chemotactic bacteria in the context of population growth, Bull. of Math. Biol. 46 (1984), pp. 19–40.
- [80] D. Le and H. L. Smith, Steady states of models of microbial growth and competition with chemotaxis, J. M. A. A. 229 (1999), pp. 295–318.
- [81] H. Levine and B. Sleeman A system of reaction diffusion equations arising in the theory of reinforced random walks, SIAM J. Appl. Math 57 (1997), pp. 683–730.
- [82] Y. Li and I. Shafrir, Blow-up analysis for solutions of  $-\Delta u = Ve^u$  in dimension two, Ind. Univ. Math. J. 43 (1994), pp. 1255–1270.
- [83] C.-S. Lin, W.-M. Ni and I. Takagi, Large amplitude stationary solutions to a chemotaxis system, J. Differ. Equations, 72 (1988), pp. 1–27.
- [84] J.-L. Martiel and A. Goldbeter, A model based on receptor desensitization for cyclic AMP signaling in Dictyostelium cells, Biophys. J. 52 (1987), pp. 807–828.
- [85] Y. Mizutani and T. Nagai, Self-similar radial solutions to a system of partial differential equations modelling chemotaxis. Bull. Kyushu Inst. Tech. Math. Natur. Sci. 42 (1995), pp. 19–28.
- [86] Y. Mizutani, N. Muramoto and K. Yoshida, Self-similar radial solutions to a parabolic system modelling chemotaxis via variational method, Hiroshima Math. J. 29 (1999), pp. 145–160.
- [87] N. Muramoto, Y. Naito and K. Yoshida, Existence of self-similar solutions to a parabolic system modeling chemotaxis, Jpn. J. Ind. Appl. Math. 17 (2000), pp. 427–451.
- [88] J. D. Murray, Modelling the pattern generating mechanism in the formation of stripes on alligators, Mathematical physics, 9th Int. Congr. Swansea/UK 1988 (1989), pp. 208–213.
- [89] J. D. Murray, *Mathematical Biology I: An Introduction*, 3rd ed., Series: Interdisciplinary Applied Mathematics. Volume. 17, Springer Verlag, Heidelberg, New York, 2002.
- [90] J. D. Murray, Mathematical Biology II: Spatial Models and Biomedical Applications, 3rd ed., Series: Interdisciplinary Applied Mathematics. Volume. 18, Springer Verlag, Heidelberg, New York, 2003.
- [91] T. Nagai and T. Ikeda, *Traveling waves in a chemotactic model*, J. Math. Biol. **30** (1991), pp. 169–184.
- [92] T. Nagai, Blow-up of radially symmetric solutions to a chemotaxis system, Adv. Math. Sci. Appl. 5 (1995), pp. 581–601.
- [93] T. Nagai and T. Senba, Behavior of radially symmetric solutions of a system related to chemotaxis, Nonlinear Anal., Theory Methods Appl. 30 (1997), pp. 3837–3842.
- [94] T. Nagai, T. Senba and K. Yoshida, Application of the Moser-Trudinger inequality to a parabolic system of chemotaxis, Funkc. Ekvacioj, Ser.Int. 40 (1997), pp. 411–433.
- [95] T. Nagai, T. Senba and K. Yoshida, Global existence of solutions to the parabolic systems of chemotaxis, RIMS Kokyuroku 1009 (1997), pp. 22-28.
- [96] T. Nagai, T. Senba and T. Suzuki, *Concentration behavior of blow-up solutions for a simplified* system of chemotaxis, Preprint 1998.
- [97] T. Nagai, T. Senba and T. Suzuki, Keller-Segel system and the concentration lemma, RIMS Kokyuroku 1025 (1998), pp. 75–80.
- [98] T. Nagai and T. Senba, Global existence and blow-up of radial solutions to a parabolic-elliptic system of chemotaxis, Adv. Math. Sci. Appl. 8 (1998), pp. 145–156.

- [99] T. Nagai, Behavior of solutions to a parabolic-elliptic system modelling chemotaxis, J. Korean Math. Soc. 37 (2000), pp. 721–733.
- [100] T. Nagai, T. Senba and T. Suzuki, Chemotaxis collapse in a parabolic system of mathematical biology, Hiroshima Math. J. 30 (2000), pp. 463–497.
- [101] T. Nagai, Global existence and Blowup of solutions to a chemotaxis system, Nonlinear Analysis 47 (2001), pp. 777–787.
- [102] T. Nagai, Blowup of nonradial solutions to parabolic-elliptic systems modeling chemotaxis in two-dimensional domains, J. Inequal. Appl. 6 (2001), pp. 37–55.
- [103] Y. Naito, T. Suzuki and K. Yoshida, Self-similar solutions to a parabolic system modeling chemotaxis, J. Differential Equations 184 (2002), pp. 386–421.
- [104] E. Nakaguchi and A. Yagi, *Full discrete approximation by Galerkin Runge-Kutta methods for a parabolic system of chemotaxis*, Hokkaido Mathematical Journal **31** (2002), pp. 385–429.
- [105] E. Nakaguchi and A. Yagi, Full discrete approximations by Galerkin method for chemotaxisgrowth model, Nonlinear Anal. Theory Meth. Appl. 47 (2001), pp. 6097–6107.
- [106] V. Nanjundiah, Chemotaxis, signal relaying and aggregation morphology, J. Theor. Biol. 42 (1973), pp. 63–105.
- [107] V. Nanjundiah and S. Shweta, The determination of spatial pattern in Dictyostelium discoideum, J. Biosci. 17 (1992), pp. 353–394.
- [108] W.-M. Ni and I. Takagi, On the shape of least-energy solutions to a semilinear Neumann problem, Comm. Pure Appl. Math. 44 (1991), pp. 819–851.
- [109] W.-M. Ni and I. Takagi, Locating the peaks of least energy solutions to a semilinear Neumann problem, Duke Math. J. 70 (1993), pp. 247–281.
- [110] A. Novick-Cohen and L. A. Segel, A gradually slowing traveling band of chemotactic bacteria. J. Math. Biol. 19 (1984), pp. 125–132.
- [111] G. M. Odell, *Travelling bands of chemotactic microorganisms*, Chapter 6 of: Mathematical models in molecular and cellular biology, ed. L.A. Segel, Cambridge University Press, Cambridge (1980), p. 345.
- [112] K. Oelschläger, On the derivation of reaction-diffusion equations as limit dynamics of systems of moderately interacting stochastic many particle processes. Probab. Th. Rel. Fields 82 (1989), pp. 565–586.
- [113] K. Osaki and A. Yagi, Finite dimensional attractors for one-dimensional Keller-Segel equations. Funkcialaj Ekvacioj 44 (2001), pp. 441–469.
- [114] H. G. Othmer, S. R. Dunbar and W. Alt, Models of dispersal in biological systems. J. Math. Biol. 26 (1988), pp. 263–298.
- [115] H. G. Othmer and P. Schaap, Oscillatory cAMP Signaling in the development of Dictyostelium discoideum, Comments on Theoretical Biology 5 (1998), pp. 175–282.
- [116] H. G. Othmer and A. Stevens, Aggregation, blowup, and collapse: the ABCs of taxis in reinforced random walks. SIAM J. Appl. Math. 57 (1997), pp. 1044–1081.
- [117] H. G. Othmer and T. Hillen, The diffusion limit of transport equations. II: Chemotaxis equations, SIAM J. Appl. Math. 62 (2002), pp. 1222–1250.
- [118] C. S. Patlak, Random walk with persistence and external bias. Bull. Math: Biophys. 15 (1953), pp. 311–338.
- [119] K. Post, A non-linear parabolic system modeling chemotaxis with sensitivity functions, Dissertation, Humboldt-Universität zu Berlin, Institut für Mathematik 1999.
- [120] L. M. Prescott, J. P. Harley and D. A. Klein, *Microbiology*. Wm. C. Brown Publishers, 3rd edition: Chicago, London, 1996.
- [121] M. Rascle Sur une équation intégro-differentielle non linéaire issue de la biologie J. Diff. Eqns 32 (1979), pp. 420–453.
- [122] M. Rascle and C. Ziti, Finite time blow-up in some models of chemotaxis, J. Math. Biol. 33 (1995), pp. 388-414.
- [123] X. Ren, Least-energy solutions to a non-autonomous semilinear problem with small diffusion coefficient Electric J. Diff. Eqns 1993 (1993), pp. 1–21.
- [124] C. D. Reymond, P. Schaap, M. Véron and J. G. Williams, *Dual role of cAMP during Dic*tyostelium development, Experientia 368 (1995), pp. 1166–1174.

Übersichtsartikel

- [125] M. Rodrigo and M. Mimura, Exact solutions of a competition-diffusion system, Hiroshima Math. J. 30 (2000), pp. 257–270.
- [126] G. Rosen, On the propagation theory for bands of chemotactic bacteria, Math. Biosci. 20 (1974), pp. 185–189.
- [127] G. Rosen and S. Baloga, On the stability of steady propagating bands of chemotactic bacteria, Math. Biosci. 24 (1975), pp. 273–279.
- [128] R. Schaaf, Stationary solutions of chemotaxis systems, Trans. Am. Math. Soc. 292 (1985), pp. 531-556.
- [129] T. L. Scribner, L. A. Segel and E. H. Rogers, A numerical study of the formation and propagation of traveling bands of chemotactic bacteria, J. Theor. Biol. 46 (1974), pp. 189–219.
- [130] T. Senba, Blow-up of radially symmetric solutions to some systems of partial differential equations modelling chemotaxis, Adv. Math. Sci. Appl. 7 (1997), pp. 79–92.
- [131] T. Senba and T. Suzuki, *Chemotaxis Collapse in a parabolic-elliptic system of mathematical biology*, Preprint 1999.
- [132] T. Senba, *Behavior of solutions to a system related to chemotaxis*, Proceedings of the Equadiff 99, (eds. B. Fiedler, K. Gröger, J. Sprekels), World Scientific Publishing Co, Singapore 2000, pp. 1115–1120.
- [133] T. Senba and T. Suzuki, Some structures of stationary solutions to Keller-Segel model, N. Kenmochi (ed.), Proceedings of the international conference on free boundary problems: theory and applications, Chiba, Japan, November 7–13, 1999. I. Tokyo: Gakkotosho. GA-KUTO Int. Ser., Math. Sci. Appl. 13 (2000), pp. 282–297.
- [134] T. Senba and T. Suzuki, Some structures of the solution set for a stationary system of chemotaxis, Adv. Math. Sci. Appl. 10 (2000), pp. 191–224.
- [135] T. Senba and T. Suzuki, Local and norm behavior of blowup solutions to a parabolic system of chemotaxis, J. Korean Math. Soc. 37 (2000), pp. 929–941.
- [136] T. Senba and T. Suzuki, Behavior of solutions to a system related to chemotaxis, Nonlinear Analysis 47 (2001), pp. 2551–2562.
- [137] T. Senba and T. Suzuki, Weak solutions to a parabolic-elliptic system of chemotaxis, J. Funct. Anal. 191 (2002), pp. 17–51.
- [138] J. A. Sherratt, Chemotaxis and chemokinesis in eukaryotic cells: The Keller-Segel model equations as an approximation to a detailed model, Bulletin Math. Biol. 56 (1994), pp. 129–146.
- [139] A. Stevens, Trail following and aggregation of myxobacteria., J. Biol. Systems 3 (1995), pp. 1059–1068.
- [140] A. Stevens, Simulation of chemotaxis-equations in two space dimensions., Nonlinear Physics of complex systems - current status and future trends, J. Parisi, S. C. Müller and W. Zimmermann (eds), Springer, Berlin 1996.
- [141] A. Stevens and F. Schweitzer, Aggregartion induced by diffusing and nondiffusing media. in Dynamics of Cell and Tissue Motion, W. Alt, A. Deutsch and G. Dunn, eds. Birkhäuser, Basel, Switzerland, 1997, pp. 183–192.
- [142] A. Stevens, A Stochastic cellular automaton modeling gliding and aggregation of myxobacteria. SIAM J. Appl. Math. (electronic) 61 (2000), pp. 172–182.
- [143] A. Stevens, The derivation of chemotaxis equations as limit dynamics of moderately interacting stochastic many-particle systems., SIAM J. Appl. Math. (electronic) 61 (2000), pp. 183–212.
- [144] I. Strauss, P. D. Frymier, R. M. Ford and P. T. Cummings, Analysis of bacterial migration: II. Studies with multiple attractant gradients, AIChE J. 41 (1995), pp. 402–414.
- [145] B. D. Sleeman and H. A. Levine, Partial differential equations of chemotaxis and angiogenesis, Math. Methods Appl. Sci. 24 (2002), pp. 405–426.
- [146] M. Struwe, Variational Methods, Springer Verlag Berlin, Heidelberg, New York, second edition 1996.
- [147] M. Struwe and G. Tarantello, On multivortex solutions in Chern-Simons gauge theory, Bolletino U. M. I. 8 (1998), pp. 109–121.
- [148] O. O. Vasieva, B. N. Vasiev, V. A. Karpov and A. N. Zaikin, A model of Dictyostelium discoideum aggregation, J. theor. Biol. 171 (1994), pp. 361–367.

- [149] B. N. Vasiev, P. Hogeweg and A. V. Panfilov, Simulation of Dictyostelium discoideum aggregation via reaction-diffusion model, Physical Review Letters 73 (1994), pp. 3173–3176.
- [150] J. J. L. Velázquez, Stability of some mechanisms of chemotactic aggregation, SIAM J. Appl. Math. 62 (2002), pp. 1581–1633 (electronic).
- [151] J. J. L. Velázquez, Point dynamics in a singular limit of the Keller-Segel model Preprint of the DEPARTAMENTO DE MATEMATICA APLICADA, Universidad Complutense de Madrid 2002.
- [152] J. J. L. Velázquez, Well posedness of a model of point dynamics in a limit of the Keller-Segel model Preprint of the DEPARTAMENTO DE MATEMATICA APLICADA, Universidad Complutense de Madrid 2002.
- [153] G. Wang and J. Wei, Steady state solutions of a reaction-diffusion system modeling chemotaxis, Math. Nachr. 233/234 (2002), pp. 221–236.
- [154] X. Wang, Qualitative behavior of solutions of chemotactic diffusion systems: effects of motility and chemotaxis and dynamics, SIAM J. Math. Anal. (electronic) 31 (2000), pp. 535–560.
- [155] G. Wolansky, A critical parabolic estimate and application to nonlocal equations arising in chemotaxis, Appl. Anal. 66 (1997), pp. 291–321.
- [156] G. Wolansky, Scent and Sensitivity: Equilibria and stability of chemotactic systems in the absence of conflicts, Preprint 1998.
- [157] G. Wolansky, Multi-components chemotactic system in absence of conflicts, Euro. Jnl. of Applied Mathematics 13 (2002), pp. 641–661.
- [158] A. Yagi, Norm behavior of solutions to a parabolic system of chemotaxis, Math. Japonica 45 (1997), pp. 241–265.
- [159] Y. Yang, H. Chen and W. Liu, On existence of global solutions and blow-up to a system of reaction-diffusion equations modelling chemotaxis, SIAM J. Math. Anal. 33 (2001), pp. 763– 785.

# Mathematik erobert das Management



Ulrich Hirsch, Gunter Dueck (Hrsg.) Management by Mathematics

Erfahrungen und Erfolge von Executives und Politikern 2003. ca. 210 S. Geb. EUR 34,90 ISBN 3-528-03187-5

## DAS BUCH

Mathematik steckt in nahezu jedem Produkt und spielt eine fundamentale Rolle in sehr vielen Dienstleistungen (u.a. Finanzen, Logistik). Die Herausgeber, Prof. Gunter Dueck, Chief Technologist von IBM, und Prof. Ulrich Hirsch, Chef von Ulrich Hirsch & Partner Unternehmensberater, gehen nun einen Schritt weiter. Ihnen ist es gelungen, mehr als zwanzig Mathematikerinnen und Mathematiker in herausragenden Führungspositionen, sowohl in der Wirtschaft, als auch in der Politik, zu Beiträgen zu gewinnen. In diesen Beiträgen wird - für viele "klassische" Manager sicherlich überraschend sehr anschaulich verdeutlicht, dass Mathematik auch in vielen Managemententscheidungen steckt. Wenigstens in solchen, die gut getroffen sind!

Vieweg

Abraham-Lincoln-Straße 46 D-65189 Wiesbaden Fax 0611.78 78-420 www.vieweg.de Änderungen vorbehalten. Erhältlich beim Buchhandel oder beim Verlag.



## Jorge Ize / Alfonso Vignoli Equivariant Degree Theory

2003. xix, 361 pages. Cloth. € 98,– [D] • ISBN 3-11-017550-9 (de Gruyter Series in Nonlinear Analysis and Applications 8)

situations.

This book presents a new degree theory for maps which commute with a group of symmetries. This degree is no longer a

single integer but an element of the group of equivariant homotopy classes of maps between two spheres and depends on the orbit types of the spaces.

The authors develop the theory and applications of this degree in a self-contained presentation starting with only elementary facts. The first chapter explains the basic tools of representation theory, homotopy theory and differential equations needed in the text. Then the degree is defined

Carlo Bardaro, Julian Musielak, Gianluca Vinti Nonlinear Integral Operators and Applications

2003. xi, 201 pages. Cloth. € 88,– [D] • ISBN 3-11-017551-7 (de Gruyter Series in Nonlinear Analysis and Applications 9)

In 1903 Fredholm published his famous paper on integral equations. Since then linear integral operators have become an important tool in many areas, including the theory of Fourier series and Fourier integrals, approximation theory and summability theory, and the theory of integral and differential equations. As regards the latter, applications were soon extended beyond linear operators. In approximation theory, however, applications were limited to linear operators mainly by the fact that the notion of singularity of an integral operator was closely connected with is linearity.

This book represents the first attempt at a comprehensive treatment of approximation theory by means of nonlinear integral operators in function spaces. In particular, the fundamental notions of approximate identity for kernels of nonlinear operators and a general concept of modulus of continuity are developed in order to obtain consis-

tent approximation results. Applications to nonlinear summability, nonlinear integral equations and nonlinear sampling theory are given. In particular, the study of nonlinear sampling operators is important since the results permit the reconstruction of several classes of signals.

and its main abstract properties are derived. The next part

is devoted to the study of equivariant homotopy groups of

spheres and to the classification of equivariant maps in the

case of abelian actions. These groups are explicitely com-

puted and the effects of symmetry breaking, products and

composition are thorougly studied. The last part deals

with computations of the equivariant index of an isolated

orbit and of an isolated loop of stationary points. Here dif-

ferential equations in a variety of situations are considered: symmetry breaking, forcing, period doubling, twisted

orbits, first integrals, gradients etc. Periodic solutions of

Hamiltonian systems, in particular spring-pendulum sys-

tems, are studied as well as Hopf bifurcation for all these

GRUYTER SERIES IN NONL

Nonlinear Integral

**Operators** and

Applications

In a wider context, the material of this book represents a starting point for new areas of research in nonlinear analysis. For this reason the text is written in a style accessible not only to researchers but to advanced students as well.





de Gruyter Berlin · New York

www.deGruyter.de · Newsletter: www.deGruyter.de/Newsletter

WALTER DE GRUYTER GMBH & CO. KG · Genthiner Straße 13 · 10785 Berlin Telefon +49-(0)30-2 60 05-0 · Fax +49-(0)30-2 60 05-251 · E-Mail wdg-info@deGruyter.de

## Präzise und verständlich: Grundlagen der Quantenmechanik



#### Joachim Weidmann Lineare Operatoren in Hilberträumen Teil II: Anwendungen

2003. 404 S. (Mathematische Leitfäden, hrsg. v. Klaus D. Bierstedt, Günther Trautmann, Gottfried Köthe) Br. € 39,90 ISBN 3-519-02237-0

#### Inhalt 🖉

Spektrale Teilräume eines selbstadjungierten Operators - Sturm-Liouville-Operatoren - Eindimensionale Diracoperatoren - Periodische Differentialoperatoren - Ein-Teilchen-Schrödingeroperatoren - Separation der Variablen und Kugelflächenfunktionen - Spektraltheorie von N-Teilchen-Schrödingeroperatoren - Grundbegriffe der Streutheorie -Existenz der Wellenoperatoren - Ein eindimensionales Streuproblem

## Das Buch

Die im ersten Teil des Buchs dargestellten Grundlagen der Theorie der linearen Operatoren in Hilberträumen werden hier benutzt, um die Spektraltheorie von Ein- und Mehrteilchen-Schrödingeroperatoren sowie des Dirac-Operators eingehend zu untersuchen. Eine einfache Darstellung der Methode der Separation der Variablen und der Kugelfunktion erlaubt es, viele Operatoren durch Separation der Variablen auf einfache zurückzuführen und damit sehr detaillierte Resultate über deren Spektren zu erzielen. Die Grundlagen der "einfachen" Streutheorie, sowie deren wichtigste Resultate der letzten Jahrzehnte werden ausführlich dargestellt; abschließend werden die Grundprinzipien der Mehr-Kanal-Streuung entwickelt.





Abraham-Lincoln-Str. 46 65189 Wiesbaden Fax 0611.7878-420 www.teubner.de
### Grundlehren der mathematischen Wissenschaften







#### J. Jacod, A. N. Shiryaev

### **Limit Theorems for Stochastic Processes**

The authors, two of the international leaders in the field, propose a systematic exposition of convergence in law for stochastic processes, from the point of view of semimartingale theory, with emphasis on results that are useful for mathematical theory and mathematical statistics. This leads them to develop in detail some particularly useful parts of the general theory of stochastic processes, such as martingale problems, and absolute continuity or contiguity results. The book contains an introduction to the theory of martingales and semimartingales, random measures stochastic integrales, Skorokhod topology, etc., as well as a large number of results which have never appeared in book form, and some entirely new results.

2nd ed. 2003. XX, 661pp. (Grundlehren der mathematischen Wissenschaften, Volume 288) Hardcover € 119,95; sFr 194,00; £ 84,00 ISBN 3-540-43932-3

#### J. Martinet

### **Perfect Lattices in Euclidean Spaces**

This book discusses a beautiful and central problem in mathematics, which involves geometry, number theory, coding theory and group theory, centering on the study of extreme lattices, i.e. those on which the density attains a local maximum, and on the so-called perfection property.

Written by a leader in the field, it is closely related to, though disjoint in content from, the classic book by J.H. Conway and N.J.A. Sloane, Sphere Packings, Lattices and Groups, published in the same series as vol. 290

2003. XVIII, 523 p. (Grundlehren der mathematischen Wissenschaften, Volume 327) Hardcover € 89,95; sFr 149,50; £ 63,00 ISBN 3-540-44236-7

#### M. Van der Put, M. F. Singer Galois Theory of Linear Differential Equations

Linear differential equations form the central topic of this volume, Galois theory being the unifying theme. A large number of aspects are presented: algebraic theory especially differential Galois theory, formal theory, classification, algorithms to decide solvability in finite terms, monodromy and Hilbert's 21st problem, asymptotics and summability, the inverse problem and linear differential equations in positive characteristic.

2003. XVII, 438 p. (Grundlehren der mathematischen Wissenschaften, Volume 328) Hardcover € 89,95; sFr 149,50; £ 63,00 ISBN 3-540-44228-6

## For further information on the series **www.springer.de/math/series**

Please order from Springer • Customer Service • Haberstr. 7 • 69126 Heidelberg, Germany Tel.: +49(0)6221 • 345 • 0 • Fax: +49(0)6221 • 345 • 4229 e-mail: orders@springer.de or through your bookseller



All Euro and GBP prices are net-prices subject to local VAT, e.g. in Germany 7% VAT for books and 16% VAT for electronic products. Prices and other details are subject to change without notice. d&p · 009510x

### UNIVERSITÄT BASEL

# Assistenzprofessur (Tenure Track) in Analysis

Am Mathematischen Institut der Universität Basel ist zum 1. April 2004 eine Assistenzprofessur (Tenure Track) in Analysis zu besetzen. Bewerberinnen und Bewerber müssen in Mathematik promoviert sein. Lehrtätigkeit im Anschluss an die Promotion ist von Vorteil.

Von den Bewerberinnen und Bewerbern wird selbständige Forschungstätigkeit in Richtung Partielle Differentialgleichungen, Dynamische Systeme, Numerische Analysis oder Angewandte Analysis erwartet. Ein starkes Engagement für Lehre und Forschung auf hohem Niveau wird vorausgesetzt.

Die Universität Basel strebt eine Erhöhung des Anteils von Frauen unter den Dozierenden an. Bewerbungen von Frauen sind deshalb besonders willkommen.

Bewerbungen mit Lebenslauf, Publikationsliste, Sonderdrucken von fünf Arbeiten, Forschungsplan, Bericht über Lehrerfahrung sowie Namen und Adressen von fünf Referenzen sind bis zum 31. Oktober 2003 zu richten an:

Prof. Dr. Marcel Tanner, Dekan der Philosophisch-Naturwissenschaftlichen Fakultät der Universität Basel Klingelbergstrasse 50 CH-4056 Basel Schweiz

Kontaktadresse für zusätzliche Informationen: Prof. D. Masser Mathematisches Institut Rheinsprung 21 CH-4051 Basel Schweiz masser@math.unibas.ch





Susanne Schmitt / Horst G. Zimmer

# Elliptic Curves

2003. Approx. x, 361 pages. Hardcover. € 78,– [D] • ISBN 3-11-016808-1 (de Gruyter Studies in Mathematics 31)

The basics of the theory of elliptic curves should be

known to everybody, be he (or she) a mathematician or a computer scientist. Especially everybody concerned with cryptography should know the elements of this theory.

The purpose of the present textbook is to give an elementary introduction to elliptic curves. Since this branch of number theory is particularly accessible to computer-assisted calculations, the authors make use of it by approaching the theory under a computational point of view. Specifically, the computeralgebra package SIMATH can be applied on several occasions. However, the book can be read also by those not interested in any computations.

Of course, the theory of elliptic curves is very comprehensive and becomes correspondingly sophisticated. That is why the authors made a choice of the topics treated. Topics covered include the determination of torsion groups, computations regarding the Mordell-Weil group, height calculations, *S*-integral points. The contents is kept as elementary as possible. In this way it becomes obvious in which respect the book differs from the numerous textbooks on elliptic curves nowadays available.

### Heinz Bauer: Selecta

Edited by Herbert Heyer, Niels Jacob and Ivan Netuka

2003. xiv, 597 pages. 1 frontispiece. Hardcover. € 128,- [D] • ISBN 3-11-017350-6

Heinz Bauer (1928–2002) was one of the prominent figures in Convex Analysis and Potential Theory in the second half of the 20th century. The Bauer minimum principle and Bauer's work on Silov's boundary and the Dirichlet problem are milestones in convex analysis. Axiomatic potential theory owes him what is known by now as Bauer harmonic spaces.

These Selecta collect more than twenty of Bauer's research papers including his seminal papers in



de Gruyter <sup>Berlin · New York</sup> Convex Analysis and Porential Theory. Above his research contributions Bauer is best known for his art of writing survey articles. Five of his surveys on different topics are reprinted in this volume. Among them is the well-known article "Approximation and Abstract Boundary", for which he was awarded with the Chauvenet Price by the American Mathematical Association in 1980.

> Please order at your bookseller or directly from us Prices are subject to change



www.deGruyter.de · Newsletter: www.deGruyter.de/Newsletter

WALTER DE GRUYTER GMBH & CO. KG · Genthiner Straße 13 · 10785 Berlin Telefon +49-(0)30-2 60 05-0 · Fax +49-(0)30-2 60 05-251 · E-Mail wdg-info@deGruyter.de

### ISSN 0012-0456